



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

REGION IX

75 Hawthorne Street

San Francisco, CA 94105-3901

August 1, 1996

96 AUG 21 PM 1:01

ENVIRONMENTAL
PROTECTION AGENCYSubject: Region 9 Preliminary Remediation Goals (PRGs) 1996From: Stanford J. Smucker, Ph.D.
Regional Toxicologist (H-9-3)
Technical Support Team

To: PRG Table Mailing List

Please find the annual update to the Region 9 PRG table. The table has been revised to reflect the most current EPA toxicological and risk assessment information. Updates to EPA toxicity values were obtained from IRIS through July 1996, HEAST through May 1995, and EPA's National Center for Environmental Assessment (NCEA, formerly ECAO).

Region 9 PRGs are "evergreen" and have evolved as new methodologies and parameters have been developed. In several cases the models, equations, and assumptions presented in RAGS HHEM, *Part B, Development of Risk-Based Preliminary Remediation Goals* (1991) have been replaced with new information that is consistent with the document, *Soil Screening Guidance*, recently issued by the Office of Solid Waste and Emergency Response (OSWER), dated April 1996.

The updated PRG table also contains soil screening levels (SSLs) for protection of groundwater. The SSLs were obtained directly from EPA/OSWER's *Soil Screening Guidance* document which is available from NTIS as EPA/540/R-96/018 and EPA/540/R-95/128. Please note that because R 9 PRGs currently evaluate intermedia transfer of volatile organic chemicals (VOCs) and heavy metals from soil to air, the PRG table does not include a separate list of SSLs for the air pathway.

To help users rapidly identify substances with new PRGs, these contaminants are printed in boldface type. Changes in PRG values are either due to new toxicity constants or new physico-chemical information. This version of the table contains revised toxicity values for acetaldehyde, chlorine cyanide, 1,3-dichlorobenzene, 2-dichloroethane, endosulfan, manganese, phosphoric acid, and 1,1,1-trichloroethane. Also, 23 additional VOCs have been identified and evaluated for inhalation exposures resulting from intermedia transfer from soil and water to air.

EPA Region 9 has established a homepage on the World Wide Web which you can find at <http://www.epa.gov/region9/>. Our homepage will soon include the PRG table in downloadable form. The electronic table contains additional information not presented in the printed table (e.g. physico-chemical constants, non-cancer PRGs for carcinogens, pathway-specific PRGs, and volatilization factors for VOCs). Meanwhile, we still provide the electronic PRG table (PRG96.zip) on California Regional Water Board's BBS (510.286.0404) for those of you who have a modem.

Before relying on any number in the table, it is recommended that the user verify the numbers with an agency toxicologist or risk assessor because the toxicity / exposure information in the table may contain errors or default assumptions that need to be refined based on further evaluation. If you find an error please send me a note via email at Smucker.Stan@epamail.epa.gov or fax at 415.744.1916.

DISCLAIMER

Preliminary remediation goals (PRGs) focus on common exposure pathways and may not consider all exposure pathways encountered at CERCLA / RCRA sites (Exhibit 1-1). PRGs do not consider impact to groundwater or address ecological concerns. PRGs are specifically not intended as a (1) stand-alone decision-making tool, (2) as a substitute for EPA guidance for preparing baseline risk assessments, or (3) a rule to determine if a waste is hazardous under RCRA.

The guidance set out in this document is not final Agency action. It is not intended, nor can it be relied upon to create any rights enforceable by any party in litigation with the United States. EPA officials may decide to follow the guidance provided herein, or act at variance with the guidance, based on an analysis of specific circumstances. The Agency also reserves the right to change this guidance at any time without public notice.

1.0 INTRODUCTION

The Region 9 PRG table combines current EPA toxicity values with "standard" exposure factors to estimate contaminant concentrations in environmental media (soil, air, and water) that are protective of humans, including sensitive groups, over a lifetime. Chemical concentrations above these levels would not automatically designate a site as "dirty" or trigger a response action. However, exceeding a PRG suggests that further evaluation of the potential risks that may be posed by site contaminants is appropriate. Further evaluation may include additional sampling, consideration of ambient levels in the environment, or a reassessment of the assumptions contained in these screening-level estimates (e.g. appropriateness of route-to-route extrapolations).

The PRG concentrations presented in the table can be used to screen pollutants in environmental media, trigger further investigation, and provide an initial cleanup goal if applicable. When considering PRGs as preliminary goals, residential concentrations should be used for maximum beneficial uses of a property. Industrial concentrations are included in the table as an alternative cleanup goal for soils, but it is not recommended that industrial concentrations be used for screening sites.

Before applying PRGs as screening tools or initial goals, the user of the table should consider whether the exposure pathways and exposure scenarios at the site are fully accounted for in the PRG calculation. Region 9 PRG concentrations are based on exposure pathways for which generally accepted methods, models, and assumptions have been developed (i.e. ingestion, dermal contact, and inhalation) for specific land-use conditions and do not consider impact to groundwater or ecological receptors (see Developing a Conceptual Site Model below).

EXHIBIT 1-1
TYPICAL EXPOSURE PATHWAYS BY MEDIUM
FOR RESIDENTIAL AND INDUSTRIAL LAND USES*

EXPOSURE PATHWAYS, ASSUMING:		
MEDIUM	RESIDENTIAL LAND USE	INDUSTRIAL LAND USE
Ground Water	<i>Ingestion from drinking</i>	Ingestion from drinking
	<i>Inhalation of volatiles</i>	Inhalation of volatiles
	Dermal absorption from bathing	Dermal absorption
Surface Water	<i>Ingestion from drinking</i>	Ingestion from drinking
	<i>Inhalation of volatiles</i>	Inhalation of volatiles
	Dermal absorption from bathing	Dermal absorption
	Ingestion during swimming	
	Ingestion of contaminated fish	
Soil	<i>Ingestion</i>	<i>Ingestion</i>
	<i>Inhalation of particulates</i>	<i>Inhalation of particulates</i>
	<i>Inhalation of volatiles</i>	<i>Inhalation of volatiles</i>
	Exposure to indoor air from soil gas	Exposure to indoor air from soil gas
	Exposure to ground water contaminated by soil leachate	Exposure to ground water contaminated by soil leachate
	Ingestion via plant, meat, or dairy products	Inhalation of particulates from trucks and heavy equipment
	<i>Dermal absorption</i>	<i>Dermal absorption</i>

Footnote:

*Exposure pathways considered in the PRG calculations are indicated in boldface italics.

2.0 READING THE PRG TABLE

2.1 General Considerations

With the exceptions described below, PRGs are chemical concentrations that correspond to fixed levels of risk (i.e. either a one-in-one million [10^{-6}] cancer risk or a noncarcinogenic hazard quotient of one, whichever occurs at a lower concentration) in soil, air, and water. In most cases, where a substance causes both cancer and noncancer or systemic effects, the 10^{-6} cancer risk will result in a more stringent criteria and consequently this value is presented in the table. PRG concentrations based on cancer risk are indicated by "ca". PRG concentrations based on noncarcinogenic health threats are indicated by "nc".

If the risk-based concentrations are to be used to screen sites, it is recommended that both cancer and noncancer-based PRGs be obtained even though the printed list contains only the more restrictive of the two values. To obtain additional values (e.g. noncancer PRGs for a carcinogenic substance), the user has the following options:

- download an electronic copy of the PRG table from EPA Region 9's homepage at [http://www.epa.gov/region 9/](http://www.epa.gov/region_9/)
- download the file (PRG96.zip) from California Regional Water Board's BBS at 510.286.0404
- use the equations presented in Section 4.0 to calculate additional PRG values

It has come to my attention that some users have been multiplying the cancer PRG concentrations by 10 or 100 to set "action levels" for triggering remediation or to set less stringent cleanup levels for a specific site after considering non-risk-based factors such as (ambient levels, detection limits, or technological feasibility). This practice recognizes that there may be a range of values that may be "acceptable" for carcinogenic risk (EPA's cancer risk range is from 10^{-6} to 10^{-4}). However, this practice could lead one to overlook serious noncancer health threats and it is strongly recommended that the user consult with a toxicologist or Regional risk assessor before doing this. For carcinogens, I have indicated by asterisk ("ca*") in the PRG table where the noncancer PRGs would be exceeded if the cancer value that is listed is multiplied by 100. Two stars ("ca**") indicate that the noncancer values would be exceeded if the cancer PRG were multiplied by 10. There is no range of "acceptable" noncarcinogenic "risk" so that under no circumstances should noncancer PRGs be multiplied by 10 or 100, when setting final cleanup criteria.

In general, PRG concentrations in the table are risk-based but for soil there are two important exceptions: (1) for several volatile chemicals, PRGs are based on the soil saturation equation ("sat") and (2) for relatively less toxic inorganic and semivolatile contaminants, a non-risk based "ceiling limit" concentration is given as 10^{+5} mg/kg ("max"). PRG concentrations that are not risk-based (i.e. either "sat" or "max") should be segregated before screening multiple

pollutant risks.

In addition to Region 9 PRGs, the PRG table also includes California EPA PRGs ("CAL-Modified PRGs") for specific chemicals where CAL-EPA values may be more restrictive than the federal values; and, soil screening levels (SSLs) for protection of groundwater (see Section 2.3 below).

2.2 Toxicity Values

EPA toxicity values, known as noncarcinogenic reference doses (RfD) and carcinogenic slope factors (SF) were obtained from IRIS through July 1996, HEAST through May 1995, and EPA's National Center for Environmental Assessment (NCEA, formerly ECAO). The priority among sources of toxicological constants used are as follows: (1) IRIS (indicated by "i"), (2) HEAST ("h"), (3) NCEA ("n"), and (4) withdrawn from IRIS or HEAST and under review ("x").

Route-to-route extrapolations ("r") were frequently used when there were no toxicity values available for a given route of exposure. Oral cancer slope factors ("SF_O") and reference doses ("RfD_O") were used for both oral and inhaled exposures for organic compounds lacking inhalation values. Inhalation slope factors ("SF_i") and inhalation reference doses ("RfD_i") were used for both inhaled and oral exposures for organic compounds lacking oral values. An additional route extrapolation is the use of oral toxicity values for evaluating dermal exposures. **Although route-to-route methods are a useful screening procedure, the appropriateness of these default assumptions for specific contaminants should be verified by a toxicologist.**

To help users rapidly identify substances with new PRGs, these contaminants are printed in boldface type. This version of the table contains revised toxicity values for acetaldehyde, chlorine cyanide, 1,3-dichlorobenzene, 2-dichloroethane, endosulfan, manganese, phosphoric acid, and 1,1,1-trichloroethane.

2.3 Soil Screening Levels

Generic soil screening levels (SSLs) for the protection of groundwater have been included in the PRG table for 100 of the most common contaminants at Superfund sites. Generic SSLs are derived using default values in standardized equations presented in *Soil Screening Guidance* (available from NTIS as document numbers PB96-963502 and PB96-963505 or EPA/540/R-95/128 and EPA/540/R-96/018).

The SSLs were developed using a default dilution-attenuation factor (DAF) of 20 to account for natural processes that reduce contaminant concentrations in the subsurface. Also included are generic SSLs that assume no dilution or attenuation between the source and the receptor well (i.e., a DAF of 1). These values can be used at sites where little or no dilution or attenuation of soil leachate concentrations is expected at a site (e.g., sites with shallow water

tables, fractured media, karst topography, or source size greater than 30 acres).

Generally, if an SSL is not exceeded for the migration to groundwater pathway, the user may eliminate this pathway from further investigation.

2.4 Miscellaneous

Volatile organic chemicals (VOCs) are indicated by "1" in the VOC column of the table and are defined as those chemicals having a Henry's Law constant greater than 10^{-5} (atm-m³/mol) and a molecular weight less than 200 g/mole). These contaminants are evaluated for potential volatilization from soil / water to air using volatilization factors (see Section 4.1). Since the 1995 PRG table, 23 additional VOCs have been identified and evaluated for inhalation exposures resulting from intermedia transfer from soil and water to air. These are indicated in boldface type.

Chemical-specific dermal absorption values for contaminants in soil and dust are presented for arsenic, cadmium, pentachlorophenol, PCBs, and dioxin as recommended by EPA's Office of Research and Development (USEPA 1992a). Otherwise, default skin absorption fractions are assumed to be 0.01 and 0.10, for inorganics and organics, respectively.

3.0 USING THE PRG TABLE

The decision to use PRGs at a site will be driven by the potential benefits of having generic risk-based concentrations in the absence of site-specific risk assessments. The original intended use of PRGs was to provide initial cleanup goals for individual chemicals given specific medium and land-use combinations (see RAGS Part B, 1991), however risk-based PRGs actually have several uses in addition to providing initial goals. These include:

- Screening sites to determine further evaluation
- Prioritizing areas of concern at megasites (e.g. federal facilities)
- Calculating risks associated with multiple contaminants

A few basic procedures are recommended for using PRGs properly. These are briefly described below. Potential problems with the use of PRGs are also identified.

3.1 Developing a Conceptual Site Model

The primary condition for use of PRGs is that exposure pathways of concern and conditions at the site match those taken into account by the PRG framework. Thus, it is always necessary to develop a conceptual site model (CSM) to identify likely contaminant source areas, exposure pathways, and potential receptors. This information can be used to determine the applicability of PRGs at the site and the need for additional information. For those pathways not covered by PRGs, a risk assessment specific to these additional pathways may be necessary. Nonetheless, the PRG lookup values will still be useful in such situations for focusing further investigative efforts on the exposure pathways not addressed.

To develop a site-specific CSM, perform an extensive records search and compile existing data (e.g. available site sampling data, historical records, aerial photographs, and hydrogeologic information). Once this information is obtained, CSM worksheets such as those provided in ASTM's *Standard Guide for Risk-Based Corrective Action Applied at Petroleum Release Sites* (1995) can be used to tailor the generic worksheet model to a site-specific CSM. The final CSM diagram represents linkages among contaminant sources, release mechanisms, exposure pathways and routes and receptors. It summarizes our understanding of the contamination problem.

As a final check, the CSM should answer the following questions:

- Are there potential ecological concerns?
- Is there potential for land use other than those covered by the PRGs (that is, residential and industrial)?
- Are there other likely human exposure pathways that were not considered in development of the PRGs (e.g. impact to groundwater, local fish consumption; raising beef, dairy, or other livestock)?
- Are there unusual site conditions (e.g. large areas of contamination, high fugitive dust levels, potential for indoor air contamination)?

If any of these four conditions exist, the PRG may need to be adjusted to reflect this new information. Suggested references for evaluating pathways not currently evaluated by Region 9 PRG's are presented in Exhibit 3-1.

EXHIBIT 3-1
**SUGGESTED READINGS FOR EVALUATING SOIL CONTAMINANT
 PATHWAYS NOT CURRENTLY ADDRESSED BY REGION 9 PRGs**

EXPOSURE PATHWAY	REFERENCE
Migration of contaminants to an underlying potable aquifer	<i>Soil Screening Guidance: User's Guide and Technical Background Document</i> (USEPA 1996a,b)
Ingestion via plant uptake	<i>Technical Support Document for Land Application of Sewage Sludge</i> (USEPA 1992a)
Ingestion via meat or dairy products	<i>Estimating Exposure to Dioxin-Like Compounds - Review Draft</i> (1994a)
Inhalation of volatiles that have migrated into basements	<i>Soil Screening Guidance: User's Guide and Technical Background Document</i> (USEPA 1996a,b)
Terrestrial environmental pathways	<i>Role of the Ecological Risk Assessment in the Baseline Risk Assessment</i> (USEPA 1994b)

3.2 Background Levels Evaluation

A necessary step in determining the usefulness of Region 9 PRGs is the consideration of background contaminant concentrations. EPA may be concerned with two types of background at sites: naturally occurring and anthropogenic. Natural background is usually limited to metals whereas anthropogenic (i.e. human-made) background includes both organic and inorganic contaminants. Before embarking on an extensive sampling and analysis program to determine local background concentrations in the area, one should first compile existing data on the subject. Far too often there is pertinent information in the literature that gets ignored, resulting in needless expenditures of time and money.

Generally EPA does not clean up below natural background. If natural background concentrations are higher than the PRGs, the generic PRGs may not be the best tool for site decisionmaking. Or, an adjustment of the PRG may be needed. For example, naturally occurring arsenic frequently is higher than the soil PRG set equal to a one-in-one-million cancer risk (the point of departure), thus an alternative PRG for arsenic is provided in the lookup tables based on non-cancer endpoints that is still protective of cancer risks as well (i.e. falls within EPA's "acceptable" risk range). Because of the problems associated with adjusting PRGs to an alternate risk level, this procedure is not recommended without first

consulting a staff toxicologist at state and / or federal regulatory agencies.

Where anthropogenic background levels exceed PRGs and EPA has determined that a response action is necessary and feasible, EPA's goal will be to develop a comprehensive response to the widespread contamination. This will often require coordination with different authorities that have jurisdiction over the sources of contamination in the area.

3.3 Risk Screening

A suggested stepwise approach for screening sites with PRGs is as follows:

- Perform an extensive records search and compile existing data.
- Identify site contaminants in the PRG Table. Record the PRG concentrations for various media and note whether PRG is based on cancer risk (indicated by "ca") or noncancer hazard (indicated by "nc"). Segregate cancer PRGs from non-cancer PRGs and exclude (but don't eliminate) non-risk based PRGs ("sat" or "max").
- For cancer risk estimates, take the site-specific concentration (maximum or 95 UCL) and divide by the PRG concentrations that are designated for cancer evaluation ("ca"). Multiply this ratio by 10^{-6} to estimate chemical-specific risk. For multiple pollutants, simply add the risk for each chemical :

$$Risk = [(\frac{conc_x}{PRG_x}) + (\frac{conc_y}{PRG_y}) + (\frac{conc_z}{PRG_z})] \times 10^{-6}$$

- For non-cancer hazard estimates. Divide concentration term by its respective non-cancer PRG designated as "nc" and sum the ratios for multiple contaminants. [Note that carcinogens may also have an associated non-cancer PRG that is not listed in the printed copy of the table and these will also need to be obtained in order to complete the non-cancer evaluation.] The non-cancer ratio represents a hazard index (HI). A hazard index of 1 or less is generally considered safe . A ratio greater than 1 suggests further evaluation:

$$Hazard\ Index = [(\frac{conc_x}{PRG_x}) + (\frac{conc_y}{PRG_y}) + (\frac{conc_z}{PRG_z})]$$

For more information on screening site risks, the reader should contact EPA Region 9's Technical Support Team.

3.4 Potential Problems

As with any risk-based tool, the potential exists for misapplication. In most cases the root cause will be a lack of understanding of the intended use of Region 9 PRGs. In order to prevent misuse of PRGs, the following should be avoided:

- Applying PRGs to a site without adequately developing a conceptual site model that identifies relevant exposure pathways and exposure scenarios,
- Not considering background concentrations when choosing PRGs as cleanup goals,
- Use of PRGs as cleanup levels without the nine-criteria analysis specified in the National Contingency Plan (or, comparable analysis for programs outside of Superfund),
- Use of PRGs as cleanup levels without verifying numbers with a toxicologist,
- Use of antiquated PRG tables that have been superseded by more recent publications,
- Not considering the effects of additivity when screening multiple chemicals, and
- Adjusting PRGs upward by factors of 10 or 100 without consulting a toxicologist.

4.0 TECHNICAL SUPPORT DOCUMENTATION

Region 9 PRGs consider human exposure hazards to chemicals from contact with contaminated soils, air, and water. The emphasis of the PRG equations and technical discussion are aimed at developing initial goals for soils, since this is an area where few standards exist. For air and water, additional reference concentrations or standards are available for many chemicals (e.g. non-zero MCLGs, AWQC, and NAAQS) and consequently the discussion of these media are brief.

4.1 Inhalation of Volatiles and Fugitive Dusts

Agency toxicity criteria indicate that risks from exposure to some chemicals via inhalation far outweigh the risk via ingestion; therefore soil PRGs have been designed to address this pathway as well. The models used to calculate PRGs for inhalation of volatiles / particulates are updates of risk assessment methods presented in RAGS Part B (USEPA 1991a) and are consistent with the *Soil Screening Guidance: User's Guide and Technical Background*

Document (USEPA 1996a,b).

To address the soil-to-air pathways the PRG calculations incorporate volatilization factors (VF_s) for volatile contaminants and particulate emission factors (PEF) for nonvolatile contaminants. These factors relate soil contaminant concentrations to air contaminant concentrations that may be inhaled on-site. The VF_s and PEF equations can be broken into two separate models: an emission model to estimate emissions of the contaminant from the soil and a dispersion model to simulate the dispersion of the contaminant in the atmosphere.

It should be noted that the box model in RAGS Part B has been replaced with a dispersion term (Q/C) derived from a modeling exercise using meteorological data from 29 locations across the United States because the box model may not be applicable to a broad range of site types and meteorology and does not utilize state-of-the-art techniques developed for regulatory dispersion modeling. The dispersion model for both volatiles and particulates is the AREA-ST, an updated version of the Office of Air Quality Planning and Standards, Industrial Source Complex Model, ISC2. However, different Q/C terms are used in the VF_s and PEF equations. Los Angeles was selected as the 90th percentile data set for volatiles and Minneapolis was selected as the 90th percentile data set for fugitive dusts (USEPA 1996 a,b). A default source size of 0.5 acres was chosen for the PRG calculations. This is consistent with the default exposure area over which Region 9 typically averages contaminant concentrations in soils. If unusual site conditions exist such that the area source is substantially larger than the default source size assumed here, an alternative Q/C could be applied (see USEPA 1996a,b).

Volatilization Factor for Soils

Volatile chemicals, defined as those chemicals having a Henry's Law constant greater than 10^{-5} (atm-m³/mol) and a molecular weight less than 200 g/mole, were screened for inhalation exposures using a volatilization factor for soils (VF_s). Please note that VF_s 's are available in the electronic version of the PRG table.

The emission terms used in the VF_s are chemical-specific and were calculated from physical-chemical information obtained from a number of sources including *Superfund Exposure Assessment Manual* (SEAM, EPA 1988), *Subsurface Contamination Reference Guide* (EPA 1990a), *Fate and Exposure Data* (Howard 1991), and *Superfund Chemical Data Matrix* (USEPA 1994c). In those cases where Diffusivity Coefficients (Di) were not provided in existing literature, Di's were calculated using Fuller's Method described in SEAM. A surrogate term was required for some chemicals that lacked physico-chemical information. In these cases, a proxy chemical of similar structure was used that may over- or under-estimate the PRG for soils. Physico-chemical information is available in the electronic version of the PRG table. To access this information, the user should display the hidden columns in the table.

Equation 4-9 forms the basis for deriving generic soil PRGs for the inhalation pathway. The following parameters in the standardized equation can be replaced with specific site data to

develop a simple site-specific PRG

- Source area
- Average soil moisture content
- Average fraction organic carbon content
- Dry soil bulk density

The basic principle of the VF_s model is applicable only if the soil contaminant concentration is at or below soil saturation. Above this level, the model cannot predict an accurate VF_s . If the PRG calculated using VF_s was greater than the calculated "sat" (Equation 4-10), the PRG was set equal to "sat" in accordance with *Soil Screening Guidance* (USEPA 1996 a,b).

Volatilization Factor for Tap Water

For tap water, an upperbound volatilization constant (VF_w) is used that is based on all uses of household water (e.g. showering, laundering, and dish washing). Certain assumptions were made. For example, it is assumed that the volume of water used in a residence for a family of four is 720 L/day, the volume of the dwelling is 150,000 L and the air exchange rate is 0.25 air changes/hour (Andelman in RAGS Part B). Furthermore, it is assumed that the average transfer efficiency weighted by water use is 50 percent (i.e. half of the concentration of each chemical in water will be transferred into air by all water uses). Note: the range of transfer efficiencies extends from 30% for toilets to 90% for dishwashers.

Particulate Emission Factor for Soils

Inhalation of chemicals adsorbed to respirable particles (PM_{10}) were assessed using a default PEF equal to $1.316 \times 10^9 \text{ m}^3/\text{kg}$ that relates the contaminant concentration in soil with the concentration of respirable particles in the air due to fugitive dust emissions from contaminated soils. The generic PEF was derived using default values in Equation 4-11, which corresponds to a receptor point concentration of approximately 0.76 ug/m^3 . The relationship is derived by Cowherd (1985) for a rapid assessment procedure applicable to a typical hazardous waste site where the surface contamination provides a relatively continuous and constant potential for emission over an extended period of time (e.g. years). This represents an annual average emission rate based on wind erosion that should be compared with chronic health criteria; it is not appropriate for evaluating the potential for more acute exposures.

The impact of the PEF on the resultant PRG concentration (that combines soil exposure pathways for ingestion, skin contact, and inhalation) can be assessed by downloading the PRG tables and displaying the hidden columns. With the exception of specific heavy metals, the PEF does not appear to significantly affect most soil PRGs. Equation 4-11 forms the basis for deriving a generic PEF for the inhalation pathway. For more details regarding specific parameters used in the PEF model, the reader is referred to *Soil Screening Guidance: Technical Background Document* (USEPA 1996a).

Note: the generic PEF evaluates windborne emissions and does not consider dust emissions from traffic or other forms of mechanical disturbance that could lead to greater emissions than assumed here.

4.2 Dermal Contact with Contaminants in Soil

Much uncertainty surrounds the determination of hazards associated with skin contact with soils. One important data gap is the lack of EPA verified toxicity values for the dermal route. For screening purposes it is assumed that dermal toxicity values can be route-to-route extrapolated from oral values but this may not always be an appropriate assumption and should be checked.

Thus far, chemical-specific absorption values for skin have been recommended for only five chemicals (arsenic, cadmium, pentachlorophenol, PCBs, and dioxin) by EPA's Office of Research and Development. For all other chemicals, default absorption values for inorganics and organics are assumed to be 1 and 10 percent, respectively. At 10 % skin absorption, the dermal dose is estimated to equal an ingestion dose for adults, using the best estimate default values in *Dermal Exposure Assessment: Principles and Applications* (EPA 1992). At 1 % absorption, the dermal dose is estimated to be 10% of the oral dose (i.e. based on an adult ingestion rate of 100 mg per day). Note: worker and children intake rates (50 and 200 mg per day, respectively) yield somewhat different results.

$$\text{dermal dose} = \text{ingestion dose}$$

$$C_{SOIL} \times ABS \times AF \times SA = C_{SOIL} \times IR$$

$$ABS = \frac{(100\text{mg/day})}{[(0.2\text{mg/cm}^2\text{-day})(5000\text{cm}^2)]} = 0.10$$

4.3 SSLs for the Migration to Groundwater Pathway

The methodology for calculating SSLs for the migration to groundwater was developed to identify chemical concentrations in soil that have the potential to contaminate groundwater. Migration of contaminants from soil to groundwater can be envisioned as a two-stage process: (1) release of contaminant in soil leachate and (2) transport of the contaminant through the underlying soil and aquifer to a receptor well. The SSL methodology considers both of these fate and transport mechanisms.

SSLs are backcalculated from acceptable ground water concentrations (i.e. nonzero MCLGs, MCLs, or risk-based PRGs). First, the acceptable groundwater concentration is multiplied by a dilution factor to obtain a target leachate concentration. For example, if the dilution factor

is 10 and the acceptable ground water concentration is 0.05 mg/L, the target soil leachate concentration would be 0.5 mg/L. The partition equation (presented in the *Soil Screening Guidance* document) is then used to calculate the total soil concentration (i.e. SSL) corresponding to this soil leachate concentration.

The SSL methodology was designed for use during the early stages of a site evaluation when information about subsurface conditions may be limited. Because of this constraint, the methodology is based on conservative, simplifying assumptions about the release and transport of contaminants in the subsurface. For more on SSLs, and how to calculate site-specific SSLs versus generic SSLs presented in the PRG table, the reader is referred to the *Soil Screening Guidance* document (USEPA 1996a,b).

4.4 Exposure Factors

Default exposure factors were obtained primarily from RAGS Supplemental Guidance Standard Default *Exposure Factors* (OSWER Directive, 9285.6-03) dated March 25, 1,991 and more recent information from U.S. EPA's Office of Solid Waste and Emergency Response, U.S. EPA's Office of Research and Development, and California EPA's Department of Toxic Substances Control (see Exhibit 4-1).

Because contact rates may be different for children and adults, carcinogenic risks during the first 30 years of life were calculated using age-adjusted factors ("adj"). Use of age-adjusted factors are especially important for soil ingestion exposures, which are higher during childhood and decrease with age. However, for purposes of combining exposures across pathways, additional age-adjusted factors are used for inhalation and dermal exposures. These factors approximate the integrated exposure from birth until age 30 combining contact rates, body weights, and exposure durations for two age groups - small children and adults. Age-adjusted factors were obtained from RAGS PART B or developed by analogy (see derivations next page).

For soils only, noncarcinogenic contaminants are evaluated in children separately from adults. No age-adjustment factor is used in this case. The focus on children is considered protective of the higher daily intake rates of soil by children and their lower body weight. For maintaining consistency when evaluating soils, dermal and inhalation exposures are also based on childhood contact rates.

- (1) ingestion([mg•yr]/[kg•d]):

$$IFS_{adj} = \frac{ED_c \times IRS_c}{BW_c} + \frac{(ED_a - ED_c) \times IRS_a}{BW_a}$$

(2) skin contact([mg•yr]/[kg•d]):

$$SFS_{adj} = \frac{ED_c \times AF \times SA_c}{BW_c} + \frac{(ED_r - ED_c) \times AF \times SA_a}{BW_a}$$

(3) inhalation ([m³•yr]/[kg•d]):

$$InhF_{adj} = \frac{ED_c \times IRA_c}{BW_c} + \frac{(ED_r - ED_c) \times IRA_a}{BW_a}$$

4.5 PRG Equations

The equations used to calculate the PRGs for carcinogenic and noncarcinogenic contaminants are presented in Equations 4-1 through 4-8. The PRG equations update RAGS Part B equations. Briefly, PRGs are risk assessments run in reverse. The methodology backcalculates a soil, air, or water concentration level from a target risk (for carcinogens) or hazard quotient (for noncarcinogens). For completeness, the soil equations combine risks from ingestion, skin contact, and inhalation simultaneously. Note: the electronic version of the table also includes pathway-specific PRGs, should the user decide against combining specific exposure pathways; or, the user wants to identify the relative contribution of each pathway to exposure.

To calculate PRGs for volatile chemicals in soil, a chemical-specific volatilization factor is calculated per Equation 4-9. Because of its reliance on Henry's law, the VF_s model is applicable only when the contaminant concentration in soil is at or below saturation (i.e. there is no free-phase contaminant present). Soil saturation ("sat") corresponds to the contaminant concentration in soil at which the adsorptive limits of the soil particles and the solubility limits of the available soil moisture have been reached. Above this point, pure liquid-phase contaminant is expected in the soil. If the PRG calculated using VF_s was greater than the calculated sat, the PRG was set equal to sat, in accordance with *Soil Screening Guidance* (USEPA 1996 a,b). The updated equation for deriving sat is presented in Equation 4-10.

EXHIBIT 4-1
STANDARD DEFAULT FACTORS

<u>Symbol</u>	<u>Definition (units)</u>	<u>Default</u>	<u>Reference</u>
CSFo	Cancer slope factor oral (mg/kg-d) ⁻¹	—	IRIS, HEAST, or NCEA
CSFi	Cancer slope factor inhaled (mg/kg-d) ⁻¹	—	IRIS, HEAST, or NCEA
RfDo	Reference dose oral (mg/kg-d)	—	IRIS, HEAST, or NCEA
RfDi	Reference dose inhaled (mg/kg-d)	—	IRIS, HEAST, or NCEA
TR	Target cancer risk	10 ⁻⁶	—
THQ	Target hazard quotient	1	—
BWa	Body weight, adult (kg)	70	RAGS (Part A), EPA 1989 (EPA/540/1-89/002)
BWc	Body weight, child (kg)	15	Exposure Factors , EPA 1991 (OSWER No. 9285.6-03)
ATc	Averaging time - carcinogens (days)	25550	RAGS(Part A), EPA 1989 (EPA/540/1-89/002)
ATn	Averaging time - noncarcinogens (days)	ED*365	
SAa	25% Surface area, adult (cm ² /day)	5000	Dermal Assessment, EPA 1992 (EPA/600/8-91/011B)
SAc	25% Surface area, child (cm ² /day)	2000	Dermal Assessment, EPA 1992 (EPA/ 600/8-9/011B)
AF	Adherence factor (mg/cm ²)	0.2	Dermal Assessment, EPA 1992 (EPA/ 600/8-9/011B)
ABS	Skin absorption (unitless): — organics —Inorganics	0.1 0.01	PEA, Cal-EPA (DTSC, 1994) PEA, Cal-EPA (DTSC, 1994)
IRAA	Inhalation rate - adult (m ³ /day)	20	Exposure Factors , EPA 1991 (OSWER No. 9285.6-03)
IRAc	Inhalation rate - child (m ³ /day)	10	RAGS (Part A), EPA 1989 (EPA/540/1-89/002)
IRWa	Drinking water ingestion - adult (L/day)	2	RAGS(Part A), EPA 1989 (EPA/540/1-89/002)
IRWc	Drinking water ingestion - child (L/day)	1	PEA, Cal-EPA (DTSC, 1994)
IRSa	Soil ingestion - adult (mg/day)	100	Exposure Factors , EPA 1991 (OSWER No. 9285.6-03)
IRSc	Soil ingestion - child (mg/day),	200	Exposure Factors , EPA 1991 (OSWER No. 9285.6-03)
IRSo	Soil ingestion - occupational (mg/day)	50	Exposure Factors , EPA 1991 (OSWER No. 9285.6-03)
EFr	Exposure frequency - residential (d/y)	350	Exposure Factors , EPA 1991 (OSWER No. 9285.6-03)
EFo	Exposure frequency - occupational (d/y)	250	Exposure Factors , EPA 1991 (OSWER No. 9285.6-03)
EDr	Exposure duration - residential (years)	30*	Exposure Factors , EPA 1991 (OSWER No. 9285.6-03)
EDc	Exposure duration - child (years)	6	Exposure Factors , EPA 1991 (OSWER No. 9285.6-03)
EDo	Exposure duration - occupational (years)	25	Exposure Factors , EPA 1991 (OSWER No. 9285.6-03)
IFSadj	Age-adjusted factors for carcinogens: Ingestion factor, soils ([mg·yr]/[kg·d])	114	RAGS(Part B) , EPA 1991 (OSWER No. 9285.7-01B)
SFSadj	Skin contact factor, soils ([mg·yr]/[kg·d])	503	By analogy to RAGS (Part B)
InhFadj	Inhalation factor ([m ³ ·yr]/[kg·d])	11	By analogy to RAGS (Part B)
IFWadj	Ingestion factor, water ([L·yr]/[kg·d])	1.1	By analogy to RAGS (Part B)
VFW	Volatilization factor for water (L/m ³)	0.5	RAGS(Part B) , EPA 1991 (OSWER No. 9285.7-01B)
PEF	Particulate emission factor (m ³ /kg)	See below	Soil Screening Guidance (EPA 1996a,b)
VF _s	Volatilization factor for soil (m ³ /kg)	See below	Soil Screening Guidance (EPA 1996a,b)
sat	Soil saturation concentration (mg/kg)	See below	Soil Screening Guidance (EPA 1996a,b)

Footnote:

*Exposure duration for lifetime residents is assumed to be 30 years total. For carcinogens, exposures are combined for children (6 years) and adults (24 years).

PRG EQUATIONS

Soil Equations: For soils, equations were based on three exposure routes (ingestion, skin contact, and inhalation).

Equation 4-1: Combined Exposures to Carcinogenic Contaminants in Residential Soil

$$C(\text{mg/kg}) = \frac{TR \times AT_c}{EF_r [(\frac{IFS_{adj} \times CSF_o}{10^6 \text{mg/kg}}) + (\frac{SFS_{adj} \times ABS \times CSF_o}{10^6 \text{mg/kg}}) + (\frac{InhF_{adj} \times CSF_i}{VF_s})]}$$

Equation 4-2: Combined Exposures to Noncarcinogenic Contaminants in Residential Soil

$$C(\text{mg/kg}) = \frac{THQ \times BW_c \times AT_n}{EF_r \times ED_c [(\frac{1}{RfD_o} \times \frac{IRS_c}{10^6 \text{mg/kg}}) + (\frac{1}{RfD_o} \times \frac{SA_c \times AF \times ABS}{10^6 \text{mg/kg}}) + (\frac{1}{RfD_i} \times \frac{IRA_c}{VF_s})]}$$

Equation 4-3: Combined Exposures to Carcinogenic Contaminants in Industrial Soil

$$C(\text{mg/kg}) = \frac{TR \times BW_a \times AT_c}{EF_o \times ED_o [(\frac{IRS_o \times CSF_o}{10^6 \text{mg/kg}}) + (\frac{SA_a \times AF \times ABS \times CSF_o}{10^6 \text{mg/kg}}) + (\frac{IRA_a \times CSF_i}{VF_s})]}$$

Equation 4-4: Combined Exposures to Noncarcinogenic Contaminants in Industrial Soil

$$C(\text{mg/kg}) = \frac{THQ \times BW_a \times AT_n}{EF_o \times ED_o [(\frac{1}{RfD_o} \times \frac{IRS_o}{10^6 \text{mg/kg}}) + (\frac{1}{RfD_o} \times \frac{SA_a \times AF \times ABS}{10^6 \text{mg/kg}}) + (\frac{1}{RfD_i} \times \frac{IRA_a}{VF_s})]}$$

Footnote:

*Use VF_s for volatile chemicals (defined as having a Henry's Law Constant [atm·m³/mol] greater than 10⁻⁵ and a molecular weight less than 200 grams/mol) or PEF for non-volatile chemicals.

Tap Water Equations:

Equation 4-5: Ingestion and Inhalation Exposures to Carcinogenic Contaminants in Water

$$C(\text{ug/L}) = \frac{TR \times AT_c \times 1000 \text{ ug/mg}}{EF_r [(IFW_{adj} \times CSF_o) + (VF_w \times InhF_{adj} \times CSF_i)]}$$

Equation 4-6: Ingestion and Inhalation Exposures to Noncarcinogenic Contaminants in Water

$$C(\text{ug/L}) = \frac{THQ \times BW_a \times AT_n \times 1000 \text{ ug/mg}}{EF_r \times ED_r [(\frac{IRW_a}{RfD_o}) + (\frac{VF_w \times IRA_a}{RfD_i})]}$$

Air Equations:

Equation 4-7: Inhalation Exposures to Carcinogenic Contaminants in Air

$$C(\text{ug/m}^3) = \frac{TR \times AT_c \times 1000 \text{ ug/mg}}{EF_r \times InhF_{adj} \times CSF_i}$$

Equation 4-8: Inhalation Exposures to Noncarcinogenic Contaminants in Air

$$C(\text{ug/m}^3) = \frac{THQ \times RfD_i \times BW_a \times AT_n \times 1000 \text{ ug/mg}}{EF_r \times ED_r \times IRA_a}$$

SOIL-TO-AIR VOLATILIZATION FACTOR (VF_s)

Equation 4-9: Derivation of the Volatilization Factor

$$VF_s (m^3/kg) = (Q/C) \times \frac{(3.14 \times D_A \times T)^{1/2}}{(2 \times \rho_b \times D_A)} \times 10^{-4} (m^2/cm^2)$$

where:

$$D_A = \frac{[(\Theta_a^{10/3} D_i H' + \Theta_w^{10/3} D_w) / n^2]}{\rho_b K_d + \Theta_w + \Theta_a H'}$$

<u>Parameter</u>	<u>Definition (units)</u>	<u>Default</u>
VF _s	Volatilization factor (m ³ /kg)	—
D _A	Apparent diffusivity (cm ² /s)	—
Q/C	Inverse of the mean conc. at the center of a 0.5-acre square source (g/m ² -s per kg/m ³)	68.81
T	Exposure interval (s)	9.5 x 10 ⁸
ρ _b	Dry soil bulk density (g/cm ³)	1.5
Θ _a	Air filled soil porosity (L _{air} /L _{soil})	0.28 or n·Θ _w
n	Total soil porosity (L _{pore} /L _{soil})	0.43 or 1 - (ρ _b /ρ _s)
Θ _w	Water-filled soil porosity (L _{water} /L _{soil})	0.15
ρ _s	Soil particle density (g/cm ³)	2.65
D _i	Diffusivity in air (cm ² /s)	Chemical-specific
H	Henry's Law constant (atm·m ³ /mol)	Chemical-specific
H'	Dimensionless Henry's Law constant	Calculated from H by multiplying by 41 (USEPA 1991a)
D _w	Diffusivity in water (cm ² /s)	Chemical-specific
K _d	Soil-water partition coefficient (cm ³ /g) = K _{oc} f _{oc}	Chemical-specific
K _{oc}	Soil organic carbon-water partition coefficient (cm ³ /g)	Chemical-specific
f _{oc}	Fraction organic carbon in soil (g/g)	0.006 (0.6%)

SOIL SATURATION CONCENTRATION (sat)

Equation 4-10: Derivation of the Soil Saturation Limit

$$sat = \frac{S}{\rho_b} (K_d \rho_b + \Theta_w + H' \Theta_a)$$

<u>Parameter</u>	<u>Definition (units)</u>	<u>Default</u>
sat	Soil saturation concentration (mg/kg)	—
S	Solubility in water (mg/L-water)	Chemical-specific
ρ_b	Dry soil bulk density (kg/L)	1.5
n	Total soil porosity (L_{pore}/L_{soil})	0.43 or $1 - (\rho_b/\rho_s)$
ρ_s	Soil particle density (kg/L)	2.65
K_d	Soil-water partition coefficient (L/kg)	$K_{oc} \times f_{oc}$ (chemical-specific)
k_{oc}	Soil organic carbon/water partition coefficient (L/kg)	Chemical-specific
f_{oc}	Fraction organic carbon content of soil (g/g)	0.006 or site-specific
Θ_w	Water-filled soil porosity (L_{water}/L_{soil})	0.15
Θ_a	Air filled soil porosity (L_{air}/L_{soil})	0.28 or $n - \Theta_w$
w	Average soil moisture content (kg_{water}/kg_{soil} or L_{water}/kg_{soil})	0.1
H	Henry's Law constant (atm-m ³ /mol)	Chemical-specific
H'	Dimensionless Henry's Law constant	$H \times 41$, where 41 is a units conversion factor

SOIL-TO-AIR PARTICULATE EMISSION FACTOR (PEF)

Equation 4-11: Derivation of the Particulate Emission Factor

$$PEF(m^3/kg) = Q/C \times \frac{3600s/h}{0.036 \times (1-V) \times (U_m/U_t)^3 \times F(x)}$$

<u>Parameter</u>	<u>Definition (units)</u>	<u>Default</u>
PEF	Particulate emission factor (m ³ /kg)	1. 316 x 10 ⁹
Q/C	Inverse of the mean concentration at the center of a 0.5-acre-square source (g/m ² -s per kg/m ³)	90.80
V	Fraction of vegetative cover (unitless)	0.5
U _m	Mean annual windspeed (m/s)	4.69
U _t	Equivalent threshold value of windspeed at 7 m (m/s)	11.32
F(x)	Function dependent on U _m /U _t derived using Cowherd (1985) (unitless)	0.194

REFERENCES

- ASTM. 1995. *Standard Guide for Risk-Based Corrective Action Applied at Petroleum Release Sites.* Designation E 1739 - 95. Philadelphia, Pennsylvania.
- California EPA. 1994. *Preliminary Endangerment Assessment Guidance Manual.* Department of Toxic Substances Control, Sacramento, California.
- Cowherd, C., G. Muleski, P. Engelhart, and D. Gillette. 1985. *Rapid Assessment of Exposure to Particulate Emission from Surface Contamination.* EPA/600/8-85/002. Prepared for Office of Health and Environmental Assessment, U.S. Environmental Protection Agency, Washington, DC. NTIS PB85-192219 7AS.
- Howard, P.H. 1990. *Handbook of Environmental Fate and Exposure Data for Organic Chemicals.* Lewis Publishers, Chelsea Michigan.
- U.S. EPA. 1988. *Superfund Exposure Assessment Manual.* EPA/540/1-88/001. Office of Emergency and Remedial Response, Washington, DC.
- U.S. EPA. 1990a. *Subsurface Contamination Reference Guide.* EPA/540/2-90/011. Office of Emergency and Remedial Response, Washington, DC.

- U.S. EPA 1990b. *Exposure Factors Handbook*. EPA/600/8089/043. Office of Health and Environmental Assessment, Washington, DC.
- U.S. EPA. 1991a. *Risk Assessment Guidance for Superfund Volume 1: Human Health Evaluation Manual (Part B, Development of Risk-Based Preliminary Remediation Goals)*. Publication 9285.7-01B. Office of Emergency and Remedial Response, Washington, DC. NTIS PB92-963333.
- U.S. EPA. 1991b. *Human Health Evaluation Manual, Supplemental Guidance: Standard Default Exposure Factors*. Publication 9285.6-03. Office of Emergency and Remedial Response, Washington, DC. NTIS PB91-921314.
- U.S. EPA. 1992a *Technical Support Document for Land Application of Sewage Sludge; Volumes I and II*. Office of Water, Washington, DC. 822/R-93-001a,b.
- U.S. EPA. 1992b *Dermal Exposure Assessment: Principles and Applications*. EPA/600/8-91/011B. Office of Health and Environmental Assessment, Washington, DC.
- U.S. EPA 1994a. *Estimating Exposure to Dioxin-Like Compounds*. U.S. EPA Office of Research and Development, EPA/600/6-88/005B.
- U.S. EPA 1994b. *Role of Ecological Assessment in the Baseline Risk Assessment*. OSWER Directive No. 9285.7-17. Office of Solid Waste and Emergency Response, Washington, DC.
- U.S. EPA 1994c. *Superfund Chemical Data Matrix*. EPA/540/R-94/009. Office of Solid Waste and Emergency Response, Washington, DC. PB94-963506.
- U.S. EPA. 1995. *Health Effects Assessment Summary Tables (HEAST): Annual Update, FY 1994*. Environmental Criteria Assessment Office, Office of Health and Environmental Assessment, Office of Research and Development, Cincinnati, OH.
- U.S. EPA. 1996a. *Soil Screening Guidance: Technical Background Document*. EPA/540/R-95/128. Office of Emergency and Remedial Response, Washington, DC. PB96-963502.
- U.S. EPA. 1996b. *Soil Screening Guidance: User's Guide*. EPA/540/R-96/018. Office of Emergency and Remedial Response, Washington, DC. PB96-963505.
- U.S. EPA. 1996c. *Integrated Risk Information System (IRIS)*. Duluth, MN.

Key: i=IRIS h=HEAST n=NCEA x=WITHDRAWN r=ROUTE EXTRAPOLATION c=CANCER PRG nc=NONCANCER PRG sat=SOIL SATURATION max=CEILING LIMIT *(where: nc < 100X ca) **(where: nc < 10X ca)

FOR PLANNING PURPOSES**TOXICITY INFORMATION****CONTAMINANT****PRELIMINARY REMEDIAL GOALS (PRGs)****SOIL SCREENING LEVELS**

SF ₀ 1/(mg/kg-d)	RfD ₀ (mg/kg-d)	SFI 1/(mg/kg-d)	RfD ₁ (mg/kg-d)	C skin V skin O abs. C soils	CAS No.	Residential Soil (mg/kg)	Industrial Soil (mg/kg)	Ambient Air (ug/m ³)	Tap Water (ug/l)	Migration to Ground Water	DAF 20 (mg/kg)	DAF 1 (mg/kg)
				0.01	n/a							
1.0E-01 h		0 0.01	542-82-1									
4.0E-02 i		0 0.01	592-01-8									
6.0E-02 i		0 0.01	508-77-4									
5.0E-03 i		0 0.01	544-92-3									
4.0E-02 i		0 0.10	460-18-5									
9.0E-02 i		0 0.10	508-68-3									
5.0E-02 i		0 0.10	506-77-4									
2.0E-02 i		0 0.10	57-12-5									
	8.6E-04 i	1 0.10	74-90-8									
2.0E-02 i		0 0.10	151-50-8									
5.0E-01 i		0 0.10	506-61-8									
1.0E-01 i		0 0.10	506-64-9									
4.0E-02 i		0 0.10	143-33-9									
5.0E-02 i		0 0.10	657-21-1									
5.0E+00 i	5.0E+00 r	0 0.10	108-94-1									
2.0E-01 i	2.0E-01 r	0 0.10	108-91-8									
5.0E-03 i	5.0E-03 r	0 0.10	68085-85-8									
1.0E-02 i	1.0E-02 r	0 0.10	52315-07-8									
7.5E-03 i	7.5E-03 r	0 0.10	66215-27-8									
1.0E-02 i	1.0E-02 r	0 0.10	1881-32-1									
3.0E-02 i	3.0E-02 r	0 0.10	75-99-0									
2.5E-02 i	2.5E-02 r	0 0.10	39515-41-8									
2.4E-01 i	2.4E-01 r	0 0.10	72-54-8									
3.4E-01 i	3.4E-01 r	0 0.10	72-65-8									
3.4E-01 i	5.0E-04 i	5.0E-04 r	0 0.10	50-29-3								
1.0E-02 i	1.0E-02 r	0 0.10	1163-19-5									
4.0E-05 i	4.0E-05 r	0 0.10	8065-48-3									
6.1E-02 h	6.1E-02 r	0 0.10	2303-16-4									
9.0E-04 h	9.0E-04 r	0 0.10	333-41-5									
4.0E-03 n	4.0E-03 r	1 0.10	132-64-9									
1.0E-02 i	1.0E-02 r	0 0.10	106-37-8									
8.4E-02 i	8.4E-02 r	2.0E-02 r	0 0.10	124-48-1								
1.4E+00 h	5.7E-05 r	2.4E-03 h	5.7E-05 i	0 0.10	96-12-8	1,2-Dibromo-3-chloropropane "CAL-Modified PRG" (PEA, 1994)						
8.5E+01 i	5.7E-05 r	7.7E-01 i	5.7E-05 h	1 0.10	106-93-4	1,2-Dibromoethane						
1.0E-01 i	1.0E-01 r	0 0.10	84-74-2			Dibutyl phthalate						
3.0E-02 i	3.0E-02 r	0 0.10	1918-00-8			Dicamba						
9.0E-02 i	5.7E-02 h	1 0.10	95-50-1			1,2-Dichlorobenzene						
3.0E-02 s	3.0E-02 r	1 0.10	541-73-1			1,3-Dichlorobenzene						
2.4E-02 h	2.4E-02 r	2.3E-01 i	1 0.10	106-48-7		1,4-Dichlorobenzene						
4.5E-01 i	4.5E-01 r	0 0.10	91-94-1			3,3-Dichlorobenzidine						
9.3E+00 r		6.3E+00 h	1 0.10	764-41-0		1,4-Dichloro-2-butene						
						Dichlorodifluoromethane						
2.0E-01 i		5.7E-02 h	1 0.10	75-71-8		1,1-Dichloroethane						
1.0E-01 h		1.4E-01 h	1 0.10	75-34-3								
9.1E-02 i	2.9E-03 r	9.1E-02 i	1 0.10	107-08-2		1,2-Dichloroethane (EDC)						
6.0E-01 i	9.0E-03 i	1.8E-01 i	1 0.10	75-35-4		1,1-Dichloroethylene						

Key: I=IRIS H=HEAST n=NCEA x=WITHDRAWN r=ROUTE EXTRAPOLATION ca=CANCER PRG nc=NONCANCER PRG sat=SOIL SATURATION max=CEILING LIMIT *(where: nc < 100X ca) **(where: nc < 10X ca)

FOR PLANNING PURPOSES

TOXICITY INFORMATION
CONTAMINANT
PRELIMINARY REMEDIAL GOALS (PRGs)
SOIL SCREENING LEVELS

SFo 1/(mg/kg-d)	RfDo (mg/kg-d)	SFI 1/(mg/kg-d)	RfDI (mg/kg-d)	O abs. O soils	CAS No.	Residential Soil (mg/kg)	Industrial Soil (mg/kg)	Ambient Air (ug/m ³)	Tap Water (ug/l)	Migration to Ground Water DAF 20 (mg/kg)	DAF 1 (mg/kg)		
1.0E-02 h		1.0E-02 r	1	0.10	150-58-2	1,2-Dichloroethylene (cis)	3.1E+01 nc	1.0E+02 nc	3.7E+01 nc	6.1E+01 nc	4.0E-01	2.0E-02	
2.0E-02 h		2.0E-02 r	1	0.10	156-60-5	1,2-Dichloroethylene (trans)	7.8E+01 nc	2.7E+02 nc	7.3E+01 nc	1.2E+02 nc	7.0E-01	3.0E-02	
9.0E-03 h		9.0E-03 r	1	0.10	540-59-0	1,2-Dichloroethylene (mixture)	3.5E+01 nc	1.2E+02 nc	3.3E+01 nc	5.5E+01 nc			
3.0E-03 I		3.0E-03 r	0	0.10	120-83-2	2,4-Dichlorophenoil	2.0E+02 nc	2.0E+03 nc	1.1E+01 nc	1.1E+02 nc	1.0E+00	5.0E-02	
8.0E-03 I		8.0E-03 r	0	0.10	94-82-6	4-(2,4-Dichlorophenoxy)butyric Acid (2,4-DB)	5.2E+02 nc	5.5E+03 nc	2.9E+01 nc	2.9E+02 nc			
1.0E-02 I		1.0E-02 r	0	0.10	94-75-7	2,4-Dichlorophenoxyacetic Acid (2,4-D)	6.5E+02 nc	6.8E+03 nc	3.7E+01 nc	3.7E+02 nc			
6.8E-02 h	1.1E-03 r	6.8E-02 r	1.1E-03 I	1	0.10	78-87-5	1,2-Dichloropropane	3.1E-01 ca*	6.8E-01 ca*	9.9E-02 ca*	1.6E-01 ca*	3.0E-02	1.0E-03
1.8E-01 h	3.0E-04 I	1.3E-01 h	5.7E-03 I	1	0.10	542-75-6	1,3-Dichloropropene	2.5E-01 ca*	5.5E-01 ca	5.2E-02 ca	8.1E-02 ca	4.0E-03	2.0E-04
		3.0E-03 I		0	0.10	616-23-9	2,3-Dichloropropanol	2.0E+02 nc	2.0E+03 nc	1.1E+01 nc	1.1E+02 nc		
2.9E-01 I	5.0E-04 I	2.9E-01 r	1.4E-04 I	0	0.10	62-73-7	Dichlorvos	1.5E+00 ca**	6.6E+00 ca*	2.3E-02 ca*	2.3E-01 ca*		
4.4E-01 x		4.4E-01 r		0	0.10	115-32-2	Dicofol	1.0E+00 ca	4.3E+00 ca	1.5E-02 ca	1.5E-01 ca		
	3.0E-02 h		5.7E-05 h	1	0.10	77-73-6	Dicyclopentadiene			2.1E-01 nc	4.2E-01 nc		
1.6E+01 I	5.0E-05 I	1.8E+01 I	5.0E-05 r	0	0.10	60-57-1	Dieldrin	2.8E-02 ca*	1.2E-01 ca	4.2E-04 ca	4.2E-03 ca	4.0E-03	2.0E-04
5.7E-03 h		5.7E-03 x	0	0.10	112-34-5	Diethylene glycol, monobutyl ether	3.7E+02 nc	3.9E+03 nc	2.1E+01 nc	2.1E+02 nc			
2.0E+00 h		2.0E+00 r	0	0.10	111-90-0	Diethylene glycol, monoethyl ether	1.0E+05 max	1.0E+05 max	7.3E+03 nc	7.3E+04 nc			
1.1E-02 h		1.1E-02 r	0	0.10	617-84-5	Diethylformamide	7.2E+02 nc	7.5E+03 nc	4.0E+01 nc	4.0E+02 nc			
1.2E-03 I	6.0E-01 I	1.2E-03 r	6.0E-01 r	0	0.10	103-23-1	Di(2-ethylhexyl)adipate	3.7E+02 nc	1.6E+03 nc	5.6E+00 nc	5.6E+01 nc		
6.0E-01 I		8.0E-01 r		0	0.10	84-86-2	Diethyl phthalate	5.2E+04 nc	1.0E+05 max	2.9E+03 nc	2.9E+04 nc		
4.7E+03 h		4.7E+03 r		0	0.10	56-53-1	Diethylstilbestrol	9.5E-05 ca	4.1E-04 ca	1.4E-06 ca	1.4E-05 ca		
8.0E-02 I		8.0E-02 r	0	0.10	43222-48-6	Difenzoquat (Avenge)	5.2E+03 nc	5.5E+04 nc	2.9E+02 nc	2.9E+03 nc			
2.0E-02 I		2.0E-02 r	0	0.10	35367-38-5	Diflubenzuron	1.3E+03 nc	1.4E+04 nc	7.3E+01 nc	7.3E+02 nc			
1.1E+01 r		1.1E+01 I	1	0.10	75-37-6	1,1-Difluoroethane			4.2E+04 nc	6.9E+04 nc			
6.0E-02 I		6.0E-02 r	0	0.10	1445-75-6	Diisopropyl methylphosphonate	5.2E+03 nc	5.5E+04 nc	2.9E+02 nc	2.9E+03 nc			
2.0E-02 I		2.0E-02 r	0	0.10	55290-84-7	Dimethipin	1.3E+03 nc	1.4E+04 nc	7.3E+01 nc	7.3E+02 nc			
2.0E-04 I		2.0E-04 r	0	0.10	60-51-5	Dimethoate	1.3E+01 nc	1.4E+02 nc	7.3E-01 nc	7.3E+00 nc			
1.4E-02 h		1.4E-02 r	0	0.10	119-90-4	3,3'-Dimethoxybenzidine	3.2E+01 ca	1.4E+02 nc	4.8E-01 ca	4.8E+00 ca			
	5.7E-06 r	5.7E-06 x	1	0.10	124-40-3	Dimethylamine	6.5E-02 nc	2.4E-01 nc	2.1E-02 nc	3.5E-02 nc			
2.0E-03 I		2.0E-03 r	0	0.10	121-99-7	N-N-Dimethylaniline	1.3E+02 nc	1.4E+03 nc	7.3E+00 nc	7.3E+01 nc			
7.5E-01 h		7.5E-01 r	0	0.10	95-68-1	2,4-Dimethylaniline	5.9E-01 ca	2.5E+00 ca	9.0E-03 ca	9.0E-02 ca			
5.8E-01 h		5.8E-01 r	0	0.10	21438-86-4	2,4-Dimethylaniline hydrochloride	7.7E-01 ca	3.3E+00 ca	1.2E-02 ca	1.2E-01 ca			
9.2E+00 h		9.2E+00 r	0	0.10	119-93-7	3,3'-Dimethylbenzidine	4.8E-02 ca	2.1E-01 ca	7.3E-04 ca	7.3E-03 ca			
2.8E+00 x		3.5E+00 x	0	0.10	57-14-7	1,1-Dimethylhydrazine	1.7E-01 ca	7.3E-01 ca	1.9E-03 ca	2.6E-02 ca			
3.7E+01 x		3.7E+01 x	0	0.10	540-73-8	1,2-Dimethylhydrazine	1.2E-02 ca	5.2E-02 ca	1.8E-04 ca	1.8E-03 ca			
1.0E-01 h		8.6E-03 I	0	0.10	68-12-2	N,N-Dimethylformamide	6.5E+03 nc	6.8E+04 nc	3.1E+01 nc	3.7E+03 nc			
2.0E-02 I		2.0E-02 r	0	0.10	105-67-9	2,4-Dimethylphenol	1.3E+03 nc	1.4E+04 nc	7.3E+01 nc	7.3E+02 nc	9.0E+00	4.0E-01	
6.0E-04 I		6.0E-04 r	0	0.10	578-26-1	2,6-Dimethylphenol	3.9E+01 nc	4.1E+02 nc	2.2E+00 nc	2.2E+01 nc			
1.0E-03 I		1.0E-03 r	0	0.10	95-65-8	3,4-Dimethylphenol	6.5E+01 nc	6.8E+02 nc	3.7E+00 nc	3.7E+01 nc			
1.0E+01 h		1.0E+01 r	0	0.10	131-11-3	Dimethyl phthalate	1.0E+05 max	1.0E+05 max	3.7E+04 nc	3.7E+05 nc			
1.0E-01 I		1.0E-01 r	0	0.10	120-61-6	Dimethyl terephthalate	6.5E+03 nc	6.8E+04 nc	3.7E+02 nc	3.7E+03 nc			
2.0E-03 I		2.0E-03 r	0	0.10	131-69-5	4,6-Dinitro-o-cyclohexyl phenol	1.3E+02 nc	1.4E+03 nc	7.3E+00 nc	7.3E+01 nc			
1.0E-04 I		1.0E-04 r	0	0.10	99-65-0	1,3-Dinitrobenzene	6.5E+00 nc	6.8E+01 nc	3.7E-01 nc	3.7E+00 nc			
4.0E-04 h		4.0E-04 r	0	0.10	528-29-0	1,2-Dinitrobenzene	2.6E+01 nc	2.7E+02 nc	1.5E+00 nc	1.5E+01 nc			
4.0E-04 h		4.0E-04 r	0	0.10	100-25-4	1,4-Dinitrobenzene	2.6E+01 nc	2.7E+02 nc	1.5E+00 nc	1.5E+01 nc			
2.0E-03 I		2.0E-03 r	0	0.10	51-28-5	2,4-Dinitrophenol	1.3E+02 nc	1.4E+03 nc	7.3E+00 nc	7.3E+01 nc			
6.8E-01 I		6.8E-01 r	0	0.10	25321-14-6	Dinitrotoluene mixture	6.5E-01 ca	2.8E+00 ca	9.9E-03 ca	9.9E-02 ca	8.0E-04	4.0E-05	
2.0E-03 I		2.0E-03 r	0	0.10	121-14-2	2,4-Dinitrotoluene (also see Dinitrotoluene mixture)	1.3E+02 nc	1.4E+03 nc	7.3E+00 nc	7.3E+01 nc	8.0E-04	4.0E-05	
1.0E-03 h		1.0E-03 r	0	0.10	606-20-2	2,6-Dinitrotoluene (also see Dinitrotoluene mixture)	6.5E+01 nc	6.8E+02 nc	3.7E+00 nc	3.7E+01 nc	7.0E-04	3.0E-05	

Key: l=IRIS h=HEAST n=NCEA x=WITHDRAWN t=ROUTE EXTRAPOLATION ca=CANCER PRG nc=NONCANCER PRG sat=SOL SATURATION max=CEILING LIMIT * (where: nc < 100X ca) **(where: nc < 10X ca)

FOR PLANNING PURPOSES

TOXICITY INFORMATION

CONTAMINANT

PRELIMINARY REMEDIAL GOALS (PRGs)

SOIL SCREENING LEVELS

			Residential Soil (mg/kg)	Industrial Soil (mg/kg)	Ambient Air (ug/m ³)	Tap Water (ug/l)	DAF 20 (mg/kg)	DAF 1 (mg/kg)	
SF ₀ 1/(mg/kg-d)	RfDo 1/(mg/kg-d)	SFI 1/(mg/kg-d)	RfDI 1/(mg/kg-d)	O abs. C soils	CAS No.				
1.0E-03 l		1.0E-03 r	0 0.10	88-65-7	Dinoseb	6.5E+01 nc	6.8E+02 nc	3.7E+01 nc	
2.0E-02 h		2.0E-02 r	0 0.10	117-84-0	di-n-Octyl phthalate	1.3E+03 nc	1.0E+04 sat	7.3E+01 nc	
1.1E-02 l		1.1E-02 r	0 0.10	123-91-1	1,4-Dioxane	4.0E+01 ca	1.7E+02 ca	6.1E-01 ca	
3.0E-02 l		3.0E-02 r	0 0.10	957-51-7	Diphenamid	2.0E+03 nc	2.0E+04 nc	1.1E+02 nc	
2.5E-02 l		2.5E-02 r	0 0.10	122-39-4	Diphenylamine	1.6E+03 nc	1.7E+04 nc	9.1E+01 nc	
8.0E-01 l		7.7E-01 l	0 0.10	122-68-7	1,2-Diphenylhydrazine	5.6E-01 ca	2.4E+00 ca	8.4E-02 ca	
2.2E-03 l		2.2E-03 r	0 0.10	85-00-7	Diquat	1.4E+02 nc	1.5E+03 nc	8.0E+00 nc	
8.6E+00 h		8.6E+00 r	0 0.10	1937-37-7	Direct black 38	5.2E-02 ca	2.2E-01 ca	7.8E-04 ca	
8.1E+00 h		8.1E+00 r	0 0.10	2602-46-2	Direct blue 6	5.5E-02 ca	2.4E-01 ca	8.3E-04 ca	
9.3E+00 h		9.3E+00 r	0 0.10	16071-88-6	Direct brown 95	4.8E-02 ca	2.1E-01 ca	7.2E-04 ca	
4.0E-05 l		4.0E-05 r	0 0.10	298-04-4	Disulfoton	2.6E+00 nc	2.7E+01 nc	1.5E-01 nc	
1.0E-02 l		1.0E-02 r	0 0.10	505-29-3	1,4-Dithiane	6.5E+02 nc	6.8E+03 nc	3.7E+02 nc	
2.0E-03 l		2.0E-03 r	0 0.10	330-54-1	Diuron	1.3E+02 nc	1.4E+03 nc	7.3E+00 nc	
4.0E-03 l		4.0E-03 r	0 0.10	2439-10-3	Dodine	2.6E+02 nc	2.7E+03 nc	1.5E+01 nc	
6.0E-03 l		6.0E-03 r	0 0.10	115-29-7	Endosulfan	3.9E+02 nc	4.1E+03 nc	2.2E+01 nc	
2.0E-02 l		2.0E-02 r	0 0.10	145-73-3	Endothall	1.3E+03 nc	1.4E+04 nc	7.3E+01 nc	
3.0E-04 l		3.0E-04 r	0 0.10	72-20-8	Endrin	2.0E+01 nc	2.0E+02 nc	1.1E+00 nc	
9.9E-03 l	2.0E-03 h	4.2E-03 l	2.9E-04 l	1 0.10	108-88-8	Epichlorohydrin	7.5E+00 nc	2.5E+01 nc	1.0E+00 nc
5.7E-03 r		5.7E-03 l	0 0.10	106-88-7	1,2-Epoxybutane	3.7E+02 nc	3.9E+03 nc	2.1E+01 nc	
2.5E-02 l		2.5E-02 r	0 0.10	759-94-4	EPTC (S-Ethyl dipropylthiocarbamate)	1.6E+03 nc	1.7E+04 nc	9.1E+01 nc	
5.0E-03 l		5.0E-03 r	0 0.10	1887-87-0	Etephenon (2-chloroethyl phosphonic acid)	3.3E+02 nc	3.4E+03 nc	1.8E+01 nc	
5.0E-04 l		6.0E-04 r	0 0.10	563-12-2	Ethion	3.3E+01 nc	3.4E+02 nc	1.8E+00 nc	
4.0E-01 h		5.7E-02 l	0 0.10	110-80-5	2-Ethoxyethanol	2.6E+04 nc	1.0E+05 max	2.1E+02 nc	
3.0E-01 h		3.0E-01 r	0 0.10	111-15-9	2-Ethoxyethanol acetate	2.0E+04 nc	1.0E+05 max	1.1E+03 nc	
8.0E-01 l		9.0E-01 r	1 0.10	141-78-6	Ethyl acetate	1.8E+04 nc	1.0E+05 max	3.3E+03 nc	
4.8E-02 h		4.8E-02 r	1 0.10	140-88-5	Ethyl acrylate	2.1E-01 ca	4.5E-01 ca	1.4E-01 ca	
1.0E-01 l		2.9E-01 l	1 0.10	100-41-4	Ethylbenzene	2.3E+02 sat	2.3E+02 sat	1.1E+03 nc	
3.0E-01 h		3.0E-01 r	0 0.10	109-78-4	Ethylene cyanohydrin	2.0E+04 nc	1.0E+05 max	1.1E+03 nc	
2.0E-02 h		2.0E-02 r	0 0.10	107-15-3	Ethylene diamine	1.3E+03 nc	1.4E+04 nc	7.3E+02 nc	
2.0E+00 l		2.0E+00 r	0 0.10	107-21-1	Ethylene glycol	1.3E+05 nc	1.0E+05 max	7.3E+03 nc	
5.7E-03 r		5.7E-03 h	0 0.10	111-78-2	Ethylene glycol, monobutyl ether	3.7E+02 nc	3.9E+03 nc	2.1E+01 nc	
1.0E+00 h		3.5E-01 h	1 0.10	76-21-8	Ethylene oxide	1.3E-01 ca	3.2E-01 ca	1.9E-02 ca	
1.1E-01 h		6.0E-05 l	1.1E-01 r	8.0E-05 r	Ethylene thiourea (ETU)	4.0E+00 ca**	1.7E+01 ca*	6.1E-02 ca*	
2.0E-02 n		2.8E+00 l	1 0.10	75-00-3	Ethyl chloride	1.1E+03 nc	1.6E+03 sat	1.0E+04 nc	
2.0E-01 l		2.0E-01 r	1 0.10	80-29-7	Ethyl ether	1.8E+03 sat	1.8E+03 sat	7.3E+02 nc	
9.0E-02 h		9.0E-02 r	1 0.10	87-63-2	Ethyl methacrylate	1.4E+02 sat	1.4E+02 sat	3.3E+02 nc	
1.0E-05 l		1.0E-05 r	0 0.10	2104-64-5	Ethyl p-nitrophenyl phenylphosphorothioate	6.5E-01 nc	6.8E+00 nc	3.7E-02 nc	
3.0E+00 l		3.0E+00 r	0 0.10	84-72-0	Ethylphthalyl ethyl glycolate	1.0E+05 max	1.0E+05 max	1.1E+05 nc	
8.0E-03 l		8.0E-03 r	0 0.10	101200-48-0	Express	5.2E+02 nc	5.5E+03 nc	2.9E+01 nc	
2.5E-04 l		2.5E-04 r	0 0.10	22224-92-6	Fenamiphos	1.6E+01 nc	1.7E+02 nc	9.1E-01 nc	
1.3E-02 l		1.3E-02 r	0 0.10	2164-17-2	Fluometuron	8.5E+02 nc	8.9E+03 nc	4.7E+01 nc	
6.0E-02 l		6.0E-02 r	0 0.10	7782-41-4	Fluorine (soluble fluoride)	3.9E+03 nc	4.1E+04 nc	2.2E+02 nc	
8.0E-02 l		8.0E-02 r	0 0.10	59758-80-4	Fluoridone	5.2E+03 nc	5.5E+04 nc	2.9E+02 nc	
2.0E-02 l		2.0E-02 r	0 0.10	58425-91-3	Flurprimidol	1.3E+03 nc	1.4E+04 nc	7.3E+02 nc	
6.0E-02 l		6.0E-02 r	0 0.10	68332-98-5	Flutolanil	3.9E+03 nc	4.1E+04 nc	2.2E+03 nc	
1.0E-02 l		1.0E-02 r	0 0.10	69409-94-5	Flualinate	6.5E+02 nc	6.8E+03 nc	3.7E+01 nc	
3.6E-03 l	1.0E-01 l	3.5E-03 r	1.0E-01 r	0 0.10	Folpet	1.3E+02 ca*	5.5E+02 ca	1.9E+01 ca	

Key : I=IRIS h=HEAST n=NCEA x=WITHDRAWN r=ROUTE EXTRAPOLATION ca=CANCER PRG nc=NONCANCER PRG sat=SOIL SATURATION max=CEILING LIMIT *(where: nc < 100X ca) **(where: nc < 10X ca)

FOR PLANNING PURPOSES

TOXICITY INFORMATION					CONTAMINANT	PRELIMINARY REMEDIAL GOALS (PRGs)				SOIL SCREENING LEVELS			
SFo 1/(mg/kg-d)	RfDo (mg/kg-d)	SFI 1/(mg/kg-d)	RfDI (mg/kg-d)	O abs. C solid	CAS No.		Residential Soil (mg/kg)	Industrial Soil (mg/kg)	Ambient Air (ug/m ³)	Tap Water (ug/l)	Migration to Ground Water		
											DAF 20 (mg/kg)		
											DAF 1 (mg/kg)		
1.1E+00 h		1.1E+00 r	0	0.10	60-34-4	Methyl hydrazine	4.0E-01 ca	1.7E+00 ca	6.1E-03 ca	6.1E-02 ca			
8.0E-02 h		2.3E-02 h	1	0.10	106-10-1	Methyl isobutyl ketone	7.7E+02 nc	2.8E+03 nc	8.3E+01 nc	1.6E+02 nc			
8.0E-02 h		8.0E-02 r	1	0.10	80-62-6	Methyl methacrylate	7.6E+02 nc	2.8E+03 nc	2.9E+02 nc	4.9E+02 nc			
3.3E-02 h		3.3E-02 r	0	0.10	99-55-8	2-Methyl-5-nitroaniline	1.3E+01 ca	5.8E+01 ca	2.0E-01 ca	2.0E+00 ca			
2.5E-04 I		2.5E-04 r	0	0.10	298-00-0	Methyl parathion	1.6E+01 nc	1.7E+02 nc	9.1E-01 nc	9.1E+00 nc			
5.0E-02 x		5.0E-02 r	0	0.10	95-48-7	2-Methylphenol	3.3E+03 nc	3.4E+04 nc	1.8E+02 nc	1.8E+03 nc	1.5E+01	8.0E-01	
5.0E-02 x		5.0E-02 r	0	0.10	106-39-4	3-Methylphenol	3.3E+03 nc	3.4E+04 nc	1.8E+02 nc	1.8E+03 nc			
5.0E-03 h		5.0E-03 r	0	0.10	106-44-5	4-Methylphenol	3.3E+02 nc	3.4E+03 nc	1.8E+01 nc	1.8E+02 nc			
6.0E-03 h		1.1E-02 h	1	0.10	25013-15-4	Methyl styrene (mixture)	1.2E+02 nc	5.2E+02 nc	4.2E+01 nc	6.0E+01 nc			
7.0E-02 h		7.0E-02 r	1	0.10	98-83-9	Methyl styrene (alpha)	6.8E+02 sat	6.8E+02 sat	2.6E+02 nc	4.3E+02 nc			
5.0E-03 n		6.8E-01 I	1	0.10	1834-04-4	Methyl tertbutyl ether (MTBE)	9.8E+03 nc	1.0E+05 max	5.5E+02 nc	5.5E+03 nc			
1.5E-01 I		1.5E-01 r	0	0.10	51218-45-2	Metolachlor (Dual)							
2.5E-02 I		2.5E-02 r	0	0.10	21087-84-9	Metribuzin	1.6E+03 nc	1.7E+04 nc	9.1E+01 nc	9.1E+02 nc			
1.8E+00 h		1.8E+00 r	2.0E-04 r	0	0.10	2365-85-5	Mirex	2.5E-01 ca*	1.1E+00 ca	3.7E-03 ca	3.7E-02 ca		
		2.0E-03 r	0	0.10	2212-67-1	Molinate	1.3E+02 nc	1.4E+03 nc	7.3E+00 nc	7.3E+01 nc			
5.0E-03 h			0	0.01	7439-98-7	Molybdenum	3.8E+02 nc	8.5E+03 nc		1.8E+02 nc			
1.0E-01 h		1.0E-01 h	0	0.10	10599-90-3	Monochloramine	6.5E+03 nc	6.8E+04 nc	3.7E+02 nc	3.7E+03 nc			
2.0E-03 I		2.0E-03 r	0	0.10	300-76-5	Naled	1.3E+02 nc	1.4E+03 nc	7.3E+00 nc	7.3E+01 nc			
1.0E-01 I		1.0E-01 r	0	0.10	15299-99-7	Napropamide	6.5E+03 nc	6.8E+04 nc	3.7E+02 nc	3.7E+03 nc	1.3E+02	7.0E+00	
2.0E-02 I			0	0.01	7440-02-0	Nickel (soluble salts)	1.5E+03 nc	3.4E+04 nc		7.3E+02 nc			
					"CAL-Modified PRG" (PEA, 1994)	1.5E+02							
8.4E-01 I		0	0.01	n/a	Nickel refinery dust			8.0E-03 ca					
1.7E+00 I			0	0.01	12035-72-2	Nickel subsulfide		1.1E+04 ca	4.0E-03 ca				
1.5E-03 x		1.5E-03 r	0	0.10	1929-82-4	Nitrapyrin	9.8E+01 nc	1.0E+03 nc	5.5E+00 nc	5.5E+01 nc			
1.6E+00 I			0	0.10	14797-55-8	Nitrate			5.8E+04 nc				
1.0E-01 x			0	0.10	10102-43-9	Nitric Oxide	6.5E+03 nc	1.0E+05 max		3.7E+03 nc			
1.0E-01 I			0	0.10	14797-65-0	Nitrite	6.5E+03 nc	1.0E+05 max		3.7E+03 nc			
6.0E-05 r		5.7E-05 h	0	0.10	68-74-4	2-Nitroaniline	3.9E+00 nc	4.1E+01 nc	2.1E-01 nc	2.2E+00 nc			
			0	0.10	99-09-2	3-Nitroaniline							
			0	0.10	100-01-8	4-Nitroaniline							
5.0E-04 I		5.7E-04 h	1	0.10	98-95-3	Nitrobenzene	1.8E+01 nc	9.4E+01 nc	2.1E+00 nc	3.4E+00 nc	1.0E-01	7.0E-03	
7.0E-02 h		7.0E-02 r	0	0.10	67-20-9	Nitrofurantoin	4.6E+03 nc	4.8E+04 nc	2.6E+02 nc	2.6E+03 nc			
1.5E+00 h		9.4E+00 h	0	0.10	59-87-0	Nitrofurazone	3.0E-01 ca	1.3E+00 ca	7.2E-04 ca	4.5E-02 ca			
1.0E+00 x			0	0.10	101102-44-0	Nitrogen dioxide							
1.0E-01 I		1.0E-01 r	0	0.10	558-88-7	Nitroguanidine	6.5E+03 nc	6.8E+04 nc	3.7E+02 nc	3.7E+03 nc			
			0	0.10	100-02-7	4-Nitrophenol							
9.4E+00 r	5.7E-03 r	9.4E+00 h	5.7E-03 I	1	0.10	79-48-8	2-Nitropropane		7.2E-04 ca	3.5E+01 ca			
5.4E+00 I	5.6E+00 I		1	0.10	924-16-3	N-Nitrosodi-n-butylamine	2.2E-02 ca	5.5E-02 ca	1.2E-03 ca	2.0E-03 ca			
2.8E+00 I		2.8E+00 r	0	0.10	1116-54-7	N-Nitrosodieethylamine	1.6E-01 ca	6.8E-01 ca	2.4E-03 ca	2.4E-02 ca			
1.5E+02 I	1.5E+02 I		0	0.10	55-18-5	N-Nitrosodimethylamine	3.0E-03 ca	1.3E-02 ca	4.5E-05 ca	4.5E-04 ca			
5.1E+01 I	4.9E+01 I		0	0.10	62-75-9	N-Nitrosodiphenylamine	8.7E-03 ca	3.7E-02 ca	1.4E-04 ca	1.3E-03 ca			
4.9E-03 I	4.9E-03 r		0	0.10	86-30-6		9.1E+01 ca	3.9E+02 ca	1.4E+00 ca	1.4E+01 ca	1.0E+00	6.0E-02	
7.0E+00 I		7.0E+00 r	0	0.10	621-84-7	N-Nitroso di-n-propylamine	6.3E-02 ca	2.7E-01 ca	9.6E-04 ca	9.6E-03 ca	5.0E-05	2.0E-06	
2.2E+01 I	2.2E+01 r		0	0.10	10595-95-6	N-Nitroso-N-methylethylamine	2.0E-02 ca	8.7E-02 ca	3.1E-04 ca	3.1E-03 ca			
2.1E+00 I	2.1E+00 I		0	0.10	930-55-2	N-Nitrosopyrrolidine	2.1E-01 ca	9.1E-01 ca	3.1E-03 ca	3.2E-02 ca			
1.0E-02 h		1.0E-02 r	0	0.10	99-08-1	m-Nitrotoluene	6.5E+02 nc	6.8E+03 nc	3.7E+01 nc	3.7E+02 nc			
1.0E-02 h		1.0E-02 r	0	0.10	99-99-0	p-Nitrotoluene	6.5E+02 nc	6.8E+03 nc	3.7E+01 nc	3.7E+02 nc			

Key: i=IRIS h=HEAST n=NCEA x=WITHDRAWN r=ROUTE EXTRAPOLATION ca=CANCER PRG nc=NONCANCER PRG sat=SOIL SATURATION max=CEILING LIMIT *(where: nc < 100X ca) **(where: nc < 10X ca)

FOR PLANNING PURPOSES

TOXICITY INFORMATION				CONTAMINANT		PRELIMINARY REMEDIAL GOALS (PRGs)				SOIL SCREENING LEVELS			
SFo 1/(mg/kg-d)	RfDo (mg/kg-d)	SFI 1/(mg/kg-d)	RfDI (mg/kg-d)	V skin O abs.	C soils C soils	CAS No.	Residential Soil (mg/kg)	Industrial Soil (mg/kg)	Ambient Air (ug/m ³)	Tap Water (ug/l)	Migration to Ground Water DAF 20 (mg/kg)	DAF 1 (mg/kg)	
7.3E+00 i	7.3E+00 r	0 0.10	50-32-8	PAHs continued				6.1E-02 ca	2.6E-01 ca	9.2E-04 ca	9.2E-03 ca	8.0E+00	4.0E-01
7.3E-03 n	7.3E-03 r	0 0.10	218-01-9	Benzo[a]pyrene "CAL-Modified PRG" (PEA, 1994)			7.2E+00 sat	7.2E+00 sat	9.2E-01 ca	9.2E+00 ca	1.6E+02	8.0E+00	
7.3E+00 n	7.3E+00 r	0 0.10	53-70-3	Chrysene "CAL-Modified PRG" (PEA, 1994)			6.1E+00				2.0E+00	8.0E-02	
		4.0E-02 i	4.0E-02 r	Dibenz[ah]anthracene	0 0.10	208-44-0	6.1E-02 ca	2.6E-01 ca	9.2E-04 ca	9.2E-03 ca	4.3E+03	2.1E+02	
		4.0E-02 i	4.0E-02 r	Fluoranthene	1 0.10	66-73-7	2.6E+03 nc	2.7E+04 nc	1.5E+02 nc	1.5E+03 nc	5.6E+02	2.8E+01	
		7.3E-01 n	7.3E-01 r	Fluorene	0 0.10	183-39-5	9.0E+01 sat	9.0E+01 sat	1.5E+02 nc	2.4E+02 nc	1.4E+01	7.0E-01	
		4.0E-02 n	4.0E-02 r	Indeno[1,2,3-cd]pyrene	1 0.10	91-20-3	6.1E-01 ca	2.6E+00 ca	9.2E-03 ca	9.2E-02 ca	8.4E+01	4.0E+00	
		3.0E-02 i	3.0E-02 r	Naphthalene	0 0.10	129-00-0	2.4E+02 sat	2.4E+02 sat	1.5E+02 nc	2.4E+02 nc	4.2E+03	2.1E+02	
		4.0E-02 n	4.0E-02 r	Pyrene	1 0.10	87747-09-5	1.0E+02 sat	1.0E+02 sat	1.1E+02 nc	1.8E+02 nc			
		1.5E-01 i	9.0E-03 r	Prochloraz	0 0.10		3.0E+00 ca	1.3E+01 ca	4.5E-02 ca	3.3E+02 ca			
		6.0E-03 h	6.0E-03 r	Profilarulin	0 0.10	26399-36-0	3.9E+02 nc	4.1E+03 nc	2.2E+01 nc	2.2E+02 nc			
		1.5E-02 i	1.5E-02 r	Prometon	0 0.10	1610-18-0	9.8E+02 nc	1.0E+04 nc	5.5E+01 nc	5.5E+02 nc			
		4.0E-03 i	4.0E-03 r	Prometryn	0 0.10	7287-19-8	2.6E+02 nc	2.7E+03 nc	1.5E+01 nc	1.5E+02 nc			
		7.5E-02 i	7.5E-02 r	Pronamide	0 0.10	23950-58-5	4.9E+03 nc	5.1E+04 nc	2.7E+02 nc	2.7E+03 nc			
		1.3E-02 i	1.3E-02 r	Propachlor	0 0.10	1918-16-7	8.5E+02 nc	8.9E+03 nc	4.7E+01 nc	4.7E+02 nc			
		5.0E-03 i	5.0E-03 r	Propanil	0 0.10	709-98-8	3.3E+02 nc	3.4E+03 nc	1.8E+01 nc	1.8E+02 nc			
		2.0E-02 i	2.0E-02 r	Propargite	0 0.10	2312-35-6	1.3E+03 nc	1.4E+04 nc	7.3E+01 nc	7.3E+02 nc			
		2.0E-03 i	2.0E-03 r	Propargyl alcohol	0 0.10	107-19-7	1.3E+02 nc	1.4E+03 nc	7.3E+00 nc	7.3E+01 nc			
		2.0E-02 i	2.0E-02 r	Propazine	0 0.10	139-40-2	1.3E+03 nc	1.4E+04 nc	7.3E+01 nc	7.3E+02 nc			
		2.0E-02 i	2.0E-02 r	Propham	0 0.10	122-42-9	1.3E+03 nc	1.4E+04 nc	7.3E+01 nc	7.3E+02 nc			
		1.3E-02 i	1.3E-02 r	Propiconazole	0 0.10	60207-90-1	8.5E+02 nc	8.9E+03 nc	4.7E+01 nc	4.7E+02 nc			
		2.0E+01 h	2.0E+01 r	Propylene glycol	0 0.10	57-55-8	1.0E+05 max	1.0E+05 max	7.3E+04 nc	7.3E+05 nc			
		7.0E-01 h	7.0E-01 r	Propylene glycol, monoethyl ether	0 0.10	111-35-3	4.6E+04 nc	1.0E+05 max	2.6E+03 nc	2.6E+04 nc			
		7.0E-01 h	5.7E-01 i	Propylene glycol, monomethyl ether	0 0.10	107-98-2	4.6E+04 nc	1.0E+05 max	2.1E+03 nc	2.6E+04 nc			
		2.4E-01 i	8.6E-03 r	Propylene oxide	1 0.10	75-58-9			5.2E-01 ca	2.2E-01 ca			
		2.5E-01 i	2.5E-01 r	Pursuit	0 0.10	61335-77-6	1.6E+04 nc	1.0E+05 max	9.1E+02 nc	9.1E+03 nc			
		2.5E-02 i	2.5E-02 r	Pydrin	0 0.10	51630-58-1	1.6E+03 nc	1.7E+04 nc	9.1E+01 nc	9.1E+02 nc			
		1.0E-03 i	1.0E-03 r	Pyridine	0 0.10	110-80-1	6.5E+01 nc	6.8E+02 nc	3.7E+00 nc	3.7E+01 nc			
		5.0E-04 i	6.0E-04 r	Quinalphos	0 0.10	13593-03-8	3.3E+01 nc	3.4E+02 nc	1.8E+00 nc	1.8E+01 nc			
		1.2E+01 h	1.2E+01 r	Quinoline	0 0.10	91-22-5	3.7E-02 ca	1.6E-01 ca	5.6E-04 ca	5.6E-03 ca			
		1.1E-01 i	3.0E-03 r	RDX (Cyclonite)	0 0.10	121-82-4	4.0E+00 ca*	1.7E+01 ca	6.1E-02 ca	6.1E-01 ca			
		3.0E-02 i	3.0E-02 r	Resmethrin	0 0.10	10453-86-8	2.0E+03 nc	2.0E+04 nc	1.1E+02 nc	1.1E+03 nc			
		5.0E-02 h	5.0E-02 r	Ronnel	0 0.10	299-84-3	3.3E+03 nc	3.4E+04 nc	1.8E+02 nc	1.8E+03 nc			
		4.0E-03 i	4.0E-03 r	Rotenone	0 0.10	83-79-4	2.6E+02 nc	2.7E+03 nc	1.5E+01 nc	1.5E+02 nc			
		2.5E-02 i	2.5E-02 r	Savay	0 0.10	78578-05-0	1.6E+03 nc	1.7E+04 nc	9.1E+01 nc	9.1E+02 nc			
		5.0E-03 i		Selenious Acid	0 0.10	7783-00-8	3.3E+02 nc	3.4E+03 nc		1.8E+02 nc	5.0E+00	3.0E-01	
		5.0E-03 i		Selenium	0 0.01	7782-49-2	3.8E+02 nc	8.5E+03 nc					
		5.0E-03 h		Selenourea	0 0.10	630-10-4	3.3E+02 nc	3.4E+03 nc		1.8E+02 nc			
		9.0E-02 i	9.0E-02 r	Sethoxydim	0 0.10	74051-80-2	5.9E+03 nc	6.1E+04 nc	3.3E+02 nc	3.3E+03 nc	3.4E+01	2.0E+00	
		5.0E-03 i		Silver and compounds	0 0.01	7440-22-4	3.8E+02 nc	8.5E+03 nc		1.8E+02 nc			
		1.2E-01 i	2.0E-03 r	Simazine	0 0.10	122-34-9	3.7E+00 ca*	1.6E+01 ca*	5.6E-02 ca	5.6E-01 ca			
		4.0E-03 i	4.0E-03 r	Sodium azide	0 0.10	26628-22-8	2.6E+02 nc	2.7E+03 nc	1.5E+01 nc	1.5E+02 nc			
		2.7E-01 h	3.0E-02 r	Sodium diethyldithiocarbamate	0 0.10	148-18-5	1.6E+00 ca	7.1E+00 ca	2.5E-02 ca	2.5E-01 ca			
		2.0E-05 i	2.0E-05 r	Sodium fluoroacetate	0 0.10	82-74-9	1.3E+00 nc	1.4E+01 nc	7.3E-02 nc	7.3E-01 nc			
		1.0E-03 h	1.0E-03 r	Sodium metavanadate	0 0.10	13718-26-8	6.5E+01 nc	6.8E+02 nc	3.7E+00 nc	3.7E+01 nc			

Key : I=IRIS, h=HEAST, n=NCEA, x=WITHDRAWN, r=ROUTE EXTRAPOLATION, ca=CANCER PRG, nc=NONCANCER PRG, sat=SOIL SATURATION, max=CEILING LIMIT * (where: nc < 100X ca) **(where: nc < 10X ca)

FOR PLANNING PURPOSES

TOXICITY INFORMATION

CONTAMINANT

PRELIMINARY REMEDIAL GOALS (PRGs)

SOIL SCREENING LEVELS

SFo 1/(mg/kg-d)	RfD 1/(mg/kg-d)	SFI 1/(mg/kg-d)	RfD O' abs. C soils	CAS No.	Residential Soil (mg/kg)	Industrial Soil (mg/kg)	Ambient Air (ug/m³)	Tap Water (ug/L)	Migration to Ground Water DAF 20 (mg/kg)	DAF (mg/kg)	
6.0E-01 I		0 0.01		7440-24-6	Strontium, stable	4.6E+04 nc	1.0E+05 max	2.2E+04 nc			
3.0E-04 I	3.0E-04 r	0 0.10		57-24-9	Strychnine	2.0E+01 nc	2.0E+02 nc	1.1E+00 nc	1.1E+01 nc		
2.0E-01 I	2.9E-01 I	1 0.10		100-42-5	Styrene	6.8E+02 sat	6.8E+02 sat	1.1E+03 nc	1.6E+03 nc	4.0E+00 2.0E-01	
2.5E-02 I	2.5E-02 r	0 0.10		88671-89-0	Systhane	1.6E+03 nc	1.7E+04 nc	9.1E+01 nc	9.1E+02 nc		
1.5E+05 h	1.5E+05 h		0 0.03	1748-01-6	2,3,7,8-TCDD (dioxin)	3.8E-06 ca	2.4E-05 ca	4.5E-08 ca	4.5E-07 ca		
7.0E-02 I	7.0E-02 r	0 0.10		34014-18-1	Tebuthiuron	4.6E+03 nc	4.8E+04 nc	2.6E+02 nc	2.6E+03 nc		
2.0E-02 h	2.0E-02 r	0 0.10		3383-08-8	Temephos	1.3E+03 nc	1.4E+04 nc	7.3E+01 nc	7.3E+02 nc		
1.3E-02 I	1.3E-02 r	0 0.10		5902-51-2	Terbacil	8.5E+02 nc	8.9E+03 nc	4.7E+01 nc	4.7E+02 nc		
2.5E-05 h		2.5E-05 r	0 0.10	13071-78-9	Terbufos	1.6E+00 nc	1.7E+01 nc	9.1E-02 nc	9.1E-01 nc		
1.0E-03 I	1.0E-03 r	0 0.10		888-50-0	Terbutryn	6.5E+01 nc	6.8E+02 nc	3.7E+00 nc	3.7E+01 nc		
3.0E-04 I	3.0E-04 r	0 0.10		95-94-3	1,2,4,5-Tetrachlorobenzene	2.0E+01 nc	2.0E+02 nc	1.1E+00 nc	1.1E+01 nc		
2.8E-02 I	3.0E-02 I	2.6E-02 I	3.0E-02 r	1 0.10	630-20-6	1,1,1,2-Tetrachloroethane	2.4E+00 ca	5.4E+00 ca	2.6E-01 ca	4.3E-01 ca	
2.0E-01 I	2.0E-01 I		1 0.10	79-34-5	1,1,2,2-Tetrachloroethane	4.5E-01 ca	1.1E+00 ca	3.3E-02 ca	5.5E-02 ca	3.0E-03 2.0E-04	
5.2E-02 n	1.0E-02 I	2.0E-03 n	1.0E-02 r	1 0.10	127-18-4	Tetrachloroethylene (PCE) "CAL-Modified PRG" (PEA, 1994)	5.4E+00 ca	1.7E+01 ca	3.3E+00 ca	1.1E+00 ca	6.0E-02 3.0E-03
3.0E-02 I	3.0E-02 r	0 0.10		58-90-2	2,3,4,6-Tetrachlorophenol	2.0E+03 nc	2.0E+04 nc	1.1E+02 nc	1.1E+03 nc		
2.0E+01 h	2.0E+01 r		0 0.10	5218-25-1	p,a,a,a-Tetrachlorotoluene	2.2E-02 ca	9.5E-02 ca	3.4E-04 ca	3.4E-03 ca		
2.4E-02 h	3.0E-02 I	2.4E-02 r	3.0E-02 r	0 0.10	Tetrachlorovinphos	1.9E+01 ca	7.9E+01 ca	2.8E-01 ca	2.8E+00 ca		
5.0E-04 I		5.0E-04 r	0 0.10	3688-24-5	Tetraethylidithiopyrophosphate	3.3E+01 nc	3.4E+02 nc	1.8E+00 nc	1.8E+01 nc		
7.0E-05 h			0 0.01	1314-32-5	Thallic oxide	5.4E+00 nc	1.2E+02 nc		2.6E+00 nc		
9.0E-05 I			0 0.01	583-68-8	Thallium acetate	6.9E+00 nc	1.5E+02 nc		3.3E+00 nc	7.0E-01 4.0E-01	
8.0E-05 I			0 0.01	6533-73-8	Thallium carbonate	6.1E+00 nc	1.4E+02 nc		2.9E+00 nc	7.0E-01 4.0E-01	
8.0E-05 I			0 0.01	7781-12-0	Thallium chloride	6.1E+00 nc	1.4E+02 nc		2.9E+00 nc	7.0E-01 4.0E-01	
8.0E-05 I			0 0.01	10102-45-1	Thallium nitrate	6.9E+00 nc	1.5E+02 nc		3.3E+00 nc	7.0E-01 4.0E-01	
8.0E-05 x			0 0.01	12039-52-0	Thallium selenite	6.9E+00 nc	1.5E+02 nc		3.3E+00 nc	7.0E-01 4.0E-01	
8.0E-05 I			0 0.01	7446-18-6	Thallium sulfate	6.1E+00 nc	1.4E+02 nc		2.9E+00 nc	7.0E-01 4.0E-01	
1.0E-02 I		1.0E-02 r	0 0.10	28249-77-6	Thiobencarb	6.5E+02 nc	6.8E+03 nc	3.7E+01 nc	3.7E+02 nc		
3.0E-02 x	3.0E-02 r	0 0.10		3688-24-5	2-(Thiocyanomethylthio)-benzothiazole (TCMTB)	2.0E+03 nc	2.0E+04 nc	1.1E+02 nc	1.1E+03 nc		
3.0E-04 h	3.0E-04 r	0 0.10		39198-18-4	Thiofanox	2.0E+01 nc	2.0E+02 nc	1.1E+00 nc	1.1E+01 nc		
8.0E-02 I		8.0E-02 r	0 0.10	23564-05-8	Thiophanate-methyl	5.2E+03 nc	5.5E+04 nc	2.9E+02 nc	2.9E+03 nc		
5.0E-03 I		5.0E-03 r	0 0.10	137-26-8	Thiram	3.3E+02 nc	3.4E+03 nc	1.8E+01 nc	1.8E+02 nc		
6.0E-01 h			0 0.01	n/a	Tin (inorganic, see tributyltin oxide for organic tin)	4.6E+04 nc	1.0E+05 max		2.2E+04 nc		
2.0E-01 I		1.1E-01 h	1 0.10	108-88-3	Toluene	7.9E+02 nc	8.8E+02 sat	4.0E+02 nc	7.2E+02 nc	1.2E+01 6.0E-01	
3.2E+00 h	3.2E+00 r		0 0.10	95-80-7	Toluene-2,4-diamine	1.4E-01 ca	6.0E-01 ca	2.1E-03 ca	2.1E-02 ca		
6.0E-01 h	6.0E-01 r		0 0.10	95-70-5	Toluene-2,5-diamine	3.9E+04 nc	1.0E+05 max	2.2E+03 nc	2.2E+04 nc		
2.0E-01 h	2.0E-01 r		0 0.10	823-40-5	Toluene-2,6-diamine	1.3E+04 nc	1.0E+05 max	7.3E+02 nc	7.3E+03 nc		
1.0E-01 I	1.9E-01 r		0 0.10	108-49-0	p-Toluidine	2.3E+00 ca	1.0E+01 ca	3.5E-02 ca	3.5E-01 ca		
1.1E+00 I	1.1E+00 I		0 0.10	8001-35-2	Toxaphene	4.0E-01 ca	1.7E+00 ca	6.0E-03 ca	6.1E-02 ca	3.1E+01 2.0E+00	
7.5E-03 I		7.5E-03 r	0 0.10	66841-25-6	Tralomethrin	4.9E+02 nc	5.1E+03 nc	2.7E+01 nc	2.7E+02 nc		
1.3E-02 I		1.3E-02 r	0 0.10	2303-17-5	Triallate	8.5E+02 nc	8.9E+03 nc	4.7E+01 nc	4.7E+02 nc		
1.0E-02 I		1.0E-02 r	0 0.10	82097-50-5	Triasulfuron	6.5E+02 nc	6.8E+03 nc	3.7E+01 nc	3.7E+02 nc		
5.0E-03 I		5.0E-03 r	0 0.10	615-54-3	1,2,4-Tribromobenzene	3.3E+02 nc	3.4E+03 nc	1.8E+01 nc	1.8E+02 nc		
3.0E-05 I		3.0E-05 r	0 0.10	58-35-9	Tributyltin oxide (TBTO)	2.0E+00 nc	2.0E+01 nc	1.1E-01 nc	1.1E+00 nc		
3.4E-02 h	3.4E-02 r		0 0.10	634-93-5	2,4,6-Trichloroaniline	1.3E+01 ca	5.6E+01 ca	2.0E-01 ca	2.0E+00 ca		
2.9E-02 h	2.9E-02 r		0 0.10	33663-50-2	2,4,6-Trichloroaniline hydrochloride	1.5E+01 ca	6.6E+01 ca	2.3E-01 ca	2.3E+00 ca		
1.0E-02 I		5.7E-02 h	1 0.10	120-82-1	1,2,4-Trichlorobenzene	5.7E+02 nc	5.5E+03 sat	2.1E+02 nc	1.9E+02 nc	5.0E+00 3.0E-01	
3.8E-02 n		3.8E-02 x	1 0.10	71-55-8	1,1,1-Trichloroethane	1.2E+03 nc	3.0E+03 sat	1.0E+03 nc	7.9E+02 nc	2.0E+00 1.0E-01	

Key : i=IRIS h=HEAST n=NCEA x=WITHDRAWN r=ROUTE EXTRAPOLATION ca=CANCER PRG nc=NONCANCER PRG sat=SOLI SATURATION max=CEILING LIMIT *(where: nc < 100X ca) **(where: nc < 10X ca)

FOR PLANNING PURPOSES

TOXICITY INFORMATION				CONTAMINANT	PRELIMINARY REMEDIAL GOALS (PRGs)				SOIL SCREENING LEVELS			
SFr 1/(mg/kg-d)	RfD (mg/kg-d)	SFr 1/(mg/kg-d)	RfD (mg/kg-d)		V skin O abd. C soils	CAS No.	Residential Soil (mg/kg)	Industrial Soil (mg/kg)	Ambient Air (ug/m ³)	Tap Water (ug/l)	Migration to Ground Water DAF 20 (mg/kg)	DAF 1 (mg/kg)
5.7E-02 i	4.0E-03 r	5.6E-02 i	4.0E-03 r	1 0.10 79-00-5	1,1,2-Trichloroethane		6.5E-01 ca	1.5E+00 ca	1.2E-01 ca	2.0E-01 ca	2.0E-02	9.0E-04
1.1E-02 n	6.0E-03 n	6.0E-03 n	6.0E-03 r	1 0.10 79-01-6	Trichloroethylene (TCE)		3.2E+00 ca*	7.0E+00 ca*	1.1E+00 ca*	1.6E+00 ca*	6.0E-02	3.0E-03
3.0E-01 i		2.0E-01 h	1 0.10 75-69-4		Trichlorofluoromethane		3.8E+02 nc	1.3E+03 nc	7.3E+02 nc	1.3E+03 nc		
1.0E-01 i	1.0E-01 r		0 0.10 95-95-4		2,4,5-Trichlorophenol		6.5E+03 nc	6.8E+04 nc	3.7E+02 nc	3.7E+03 nc	2.7E+02	1.4E+01
1.1E-02 i	1.1E-02 i		0 0.10 88-06-2		2,4,6-Trichlorophenol		4.0E+01 ca	1.7B+02 ca	6.2E-01 ca	6.1E+00 ca	2.0E-01	8.0E-03
1.0E-02 i	1.0E-02 r		0 0.10 93-76-5		2,4,5-Trichlorophenoxyacetic Acid		6.5E+02 nc	6.8E+03 nc	3.7E+01 nc	3.7E+02 nc		
6.0E-03 i	6.0E-03 r		0 0.10 93-72-1		2-(2,4,5-Trichlorophenoxy) propionic acid		5.2E+02 nc	5.5E+03 nc	2.9E+01 nc	2.9E+02 nc		
5.0E-03 i	5.0E-03 r		1 0.10 598-77-6		1,1,2-Trichloropropane		1.5E+01 nc	5.0E+01 nc	1.8E+01 nc	3.0E+01 nc		
7.0E+00 h	6.0E-03 i	7.0E+00 r	5 0E-03 r	1 0.10 98-18-4	1,2,3-Trichloropropane		1.4E-03 ca	3.1E-03 ca	9.6E-04 ca	1.6E-03 ca		
5.0E-03 h	5.0E-03 r		1 0.10 96-19-5		1,2,3-Trichloropropene		1.1E+01 nc	3.8E+01 nc	1.8E+01 nc	3.0E+01 nc		
3.0E+01 i		8.6E+00 h	1 0.10 76-13-1		1,1,2-Trichloro-1,2,2-trifluoroethane		5.6E+03 sat	5.6E+03 sat	3.1E+04 nc	5.9E+04 nc		
3.0E-03 i	3.0E-03 r		0 0.10 58138-08-2		Tridiphane		2.0E+02 nc	2.0E+03 nc	1.1E+01 nc	1.1E+02 nc		
2.0E-03 r		2.0E-03 i	1 0.10 121-44-8		Triethylamine		2.3E+01 nc	8.4E+01 nc	7.3E+00 nc	1.2E+01 nc		
7.7E-03 i	7.5E-03 i	7.7E-03 r	7.5E-03 r	0 0.10 1582-09-8	Trifluralin		5.8E+01 ca**	2.5E+02 ca*	8.7E-01 ca*	8.7E+00 ca*		
3.7E-02 h		3.7E-02 i		0 0.10 512-56-1	Trimethyl phosphate		1.2E+01 ca	5.2E+01 ca	1.8E-01 ca	1.8E+00 ca		
5.0E-05 i	5.0E-05 r		0 0.10 99-35-4		1,3,5-Trinitrobenzene		3.3E+00 nc	3.4E+01 nc	1.8E-01 nc	1.8E+00 nc		
1.0E-02 h	1.0E-02 r		0 0.10 479-45-8		Trinitrophenylmethylnitramine		6.5E+02 nc	6.8E+03 nc	3.7E+01 nc	3.7E+02 nc		
3.0E-02 i	5.0E-04 i	3.0E-02 r	5.0E-04 r	0 0.10 118-96-7	2,4,6-Trinitrotoluene		1.5E+01 ca**	6.4E+01 ca**	2.2E-01 ca**	2.2E+00 ca**		
3.0E-03 i			0 0.01 7440-61-1		Uranium (soluble salts)							
7.0E-03 h			0 0.01 7440-62-2		Vanadium		5.4E+02 nc	1.2E+04 nc		2.6E+02 nc	6.0E+03	3.0E+02
9.0E-03 i			0 0.01 1314-62-1		Vanadium pentoxide		6.9E+02 nc	1.5E+04 nc		3.3E+02 nc	6.0E+03	3.0E+02
2.0E-02 h			0 0.01 13701-70-7		Vanadium sulfate		1.5E+03 nc	3.4E+04 nc		7.3E+02 nc	6.0E+03	3.0E+02
1.0E-03 i	1.0E-03 r		0 0.10 1928-77-7		Vernam		6.5E+01 nc	6.8E+02 nc	3.7E+00 nc	3.7E+01 nc		
2.5E-02 i	2.5E-02 r		0 0.10 50471-44-8		Vinclozolin		1.6E+03 nc	1.7E+04 nc	9.1E+01 nc	9.1E+02 nc		
1.0E+00 h		5.7E-02 i	1 0.10 108-05-4		Vinyl acetate		7.8E+02 nc	2.6E+03 nc	2.1E+02 nc	4.1E+02 nc	1.7E+02	8.0E+00
1.1E-01 r	6.6E-04 r	1.1E-01 h	6.6E-04 i	1 0.10 593-80-2	Vinyl bromide (bromoethene)		1.9E-01 ca*	4.1E-01 ca*	6.1E-02 ca*	1.0E-01 ca*		
1.9E+00 h		3.0E-01 h		1 0.10 75-01-4	Vinyl chloride		1.6E-02 ca	3.5E-02 ca	2.2E-02 ca	2.0E-02 ca	1.0E-02	7.0E-04
3.0E-04 i		3.0E-04 r	0 0.10 81-81-2		Warfarin		2.0E+01 nc	2.0E+02 nc	1.1E+00 nc	1.1E+01 nc		
2.0E+00 i		2.0E-01 x	1 0.10 108-39-3		m-Xylene		3.2E+02 sat	3.2E+02 sat	7.3E+02 nc	1.4E+03 nc	2.1E+02	1.0E+01
2.0E+00 i		2.0E-01 x	1 0.10 65-47-8		o-Xylene		3.2E+02 sat	3.2E+02 sat	7.3E+02 nc	1.4E+03 nc	1.9E+02	9.0E+00
			1 0.10 108-42-3		p-Xylene		3.2E+02 sat	3.2E+02 sat				
2.0E+00 i		2.0E-01 x	1 0.10 1330-20-7		Xylene (mixed)		3.2E+02 sat	3.2E+02 sat	7.3E+02 nc	1.4E+03 nc	2.0E+02	1.0E+01
3.0E-01 i			0 0.01 7440-86-6		Zinc		2.3E+04 nc	1.0E+05 max		1.1E+04 nc	1.2E+04	6.2E+02
3.0E-04 i			0 0.01 1314-84-7		Zinc phosphide		2.3E+01 nc	5.1E+02 nc		1.1E+01 nc		
5.0E-02 i		5.0E-02 r	0 0.10 12122-67-7		Zineb		3.3E+03 nc	3.4E+04 nc	1.8E+02 nc	1.8E+03 nc		