600 8

Tier 2 default.
Inputs Meivitt Sunds

| Mamol |
|-------|
| |

| | | Resid | ential | Commercial/ Industrial |
|---|----------------------------------|-----------------------|--------------|---------------------------|
| nput Parameters | Units | Child | Adult | Worker |
| | Soil-Specific Pa | rameters | | |
| Capillary fringe thickness | cm | | (10.1) | |
| Capillary fringe air content | cm³/cm³ | | 0.025 | |
| Capillary fringe water content | cm³/cm³ | | 0.325 | |
| Fraction organic carbon (FOC*) | g oc/g soil | =adult | 0.01 | =adult |
| Groundwater Darcy velocity | cm/yr | residential | 600 | residential |
| Groundwater mixing zone thickness | cm | | 305 | : - :: |
| Infiltration rate through the vadose zone | cm/yr | | 9 | |
| Soil bulk density | g/cm ³ | | 1.72 | |
| Soil to skin adherence factor | mg/cm ² | 0.2 | 0.2 | 0.2 |
| Total soil porosity | cm ³ /cm ³ | =adult residential | 0.35 | =adult residential |
| Vadose zone air content | cm ³ /cm ³ | | 0.2 | |
| Vadose zone water content | cm³/cm³ | | 0.1 <u>5</u> | |
| Vadose zone thickness | cm | | 232.9 | |
| Str | uctural and Clim | atic Parameters | | |
| Areal fraction of cracks in building foundation | cm²/cm² | | 0.001 | 0.001 |
| Foundation air content | cm³/cm³ | | 0.26 | =adult |
| Foundation water content | cm ³ /cm ³ | | 0.12 | residential |
| Foundation thickness | cm | | 15 | 15 |
| Lower depth of surficial soil zone | cm | | 100.0 | |
| Depth to subsurface soil sources | cm | =adult residential | 100 | |
| Depth to groundwater | cm | | 243 | =adult residential |
| Width of source area parallel to wind or groundwater flow direction | cm | | 1500 | |
| Outdoor air mixing zone height | cm | | 200 | |
| Particulate emission rate | g/cm ² -s | | 1.38E-11 | 1.38E-11 |
| Wind speed above ground surface in outdoor air mixing zone | cm/s | | 322 | =adult residential |

1630 ParkSt. Merritt Sands Tier 3

Inputs

| | | Residential | | Commercial/ Industrial |
|---|----------------------------------|-----------------------|----------|---------------------------|
| Input Parameters | Units | Child | Adult | Worker |
| | Soil-Specific Pa | arameters | | |
| Capillary fringe thickness | cm | | 4.6 | |
| Capillary fringe air content | cm ³ /cm ³ | | 0.938 | |
| Capillary fringe water content | cm³/cm³ | | 0,342 | |
| Fraction organic carbon (FOC*) | g oc/g soil | =adult | 0.01 | =adult |
| Groundwater Darcy velocity | cm/yr | residential | 1500 | residential |
| Groundwater mixing zone thickness | cm | | 260 | |
| Infiltration rate through the vadose zone | cm/yr | | 15 | |
| Soil bulk density | g/cm ³ | | 1.70 | |
| Soil to skin adherence factor | mg/cm ² | 0.2 | 0.2 | 0.2 |
| Total soil porosity | cm³/cm³ | | 0.38 | =adult residential |
| Vadose zone air content | cm ³ /cm ³ | =adult | 0.26 | |
| Vadose zone water content | cm³/cm³ | residential | 0.12 | |
| Vadose zone thickness | cm | | 240 | |
| Stru | ctural and Clima | atic Parameters | | |
| Areal fraction of cracks in building foundation | cm²/cm² | | 0.001 | 0.001 |
| Foundation air content | cm³/cm³ | | 0.26 | =adult |
| Foundation water content | cm³/cm³ | 1 | 0.12 | residential |
| Foundation thickness | cm | | 15 | 15 |
| Lower depth of surficial soil zone | cm | =adult residential | 100.0 | |
| Depth to subsurface soil sources | cm | | 91 | |
| Depth to groundwater | cm | | 240 | =adult residential |
| Width of source area parallel to wind or groundwater flow direction | cm | | 2400 | |
| Outdoor air mixing zone height | cm | | 200 | |
| Particulate emission rate | g/cm²-s | | 2.20E-10 | 1.38E-11 |
| Wind speed above ground surface in outdoor air mixing zone | cm/s | | 230 | =adult residential |

Inputs

| | | Residential | | Commercial/ Industrial |
|--|----------------------------------|-----------------------|----------|---------------------------|
| Input Parameters | Units | Child | Adult | Worker |
| | Exposure Para | meters | | |
| Averaging time for carcinogens | yr | =adult residential | 70 | =adult residential |
| Averaging time for non-carcinogens | yr | 6 | 24 | 25 |
| Averaging time for vapor flux | s | =adult residential | 9.46E+08 | 7.88E+08 |
| Body weight | kg | 15 | 70 | 70 |
| Building air volume/floor area | cm ³ /cm ² | =adult residential | 229 | 305 |
| Exposure duration | yr | 6 | 24 | 25 |
| Exposure frequency | d/yr | 350 | 350 | 250 |
| Exposure frequency to water used for recreation | d/yr | 120 | 120 | 0 |
| Exposure time to indoor air | hr/d | 24 | 24 | 9 |
| Exposure time to outdoor air | hr/d | 16 | 16 | 9 |
| Exposure time to water used for recreation | hr/d | 2 | 1.0 | 0 |
| Groundwater ingestion rate | L/d | 1 | 2 | 1 |
| Indoor air exchange rate | 1/s | =adult residential | 5.60E-04 | 1.40E-03 |
| Indoor inhalation rate | m³/d | 10 | 15 | 20 |
| Ingestion rate of water used for recreation | L/hr | 0.05 | 0.05 | 0 |
| Outdoor inhalation rate | m ³ /d | 10 | 20 | 20 |
| Skin surface area exposed to soil | cm ² | 2000 | 5000 | 5000 |
| Skin surface area exposed to water used for recreation | cm ² | 8000 | 20000 | 0 |
| Soil ingestion rate | mg/d | 200 | 100 | 50 |
| | TARGET RISK | LEVELS | | |
| Individual Excess Lifetime Cancer Risk | unitless | =adult | 1.0E-05 | 1.0E-05 |
| Hazard quotient | unitless | residential | 1.0 | 1.0 |

Inputs

| * | | Residential | | Commercial/ Industrial |
|--|----------------------------------|-----------------------|----------|---------------------------|
| Input Parameters | Units | Child | Adult | Worker |
| | Exposure Para | ameters | | |
| Averaging time for carcinogens | yr | =adult residential | 70 | =adult residential |
| Averaging time for non-carcinogens | yr | 6 | 30 | 25 |
| Averaging time for vapor flux | s | =adult residential | 9.46E+08 | 7.88E+08 |
| Body weight | kg | 15 | 70 | 70 |
| Building air volume/floor area | cm ³ /cm ² | =adult residential | 200 | 300 |
| Exposure duration | yr | 6 | 30 | 25 |
| Exposure frequency | d/yr | 350 | 350 | 180 |
| Exposure frequency to water used for recreation | d/yr | 120 | 120 | 0 |
| Exposure time to indoor air | hr/d | 24 | 24 | 9 |
| Exposure time to outdoor air | hr/d | 16 | 16 | 9 |
| Exposure time to water used for recreation | hr/d | 2 | 1.0 | 0 |
| Groundwater ingestion rate | L/d | 1 | 2 | 1 |
| Indoor air exchange rate | 1/s | =adult residential | 5.60E-04 | 1.40E-03 |
| Indoor inhalation rate | m³/d | 10 | 15 | 20 |
| Ingestion rate of water used for recreation | L/hr | 0.05 | 0.05 | 0 |
| Outdoor inhalation rate | m³/d | 10 | 20 | 10 |
| Skin surface area exposed to soil | cm ² | 2000 | 5800 | 5800 |
| Skin surface area exposed to water used for recreation | cm ² | 8000 | 20000 | 0 |
| Soil ingestion rate | mg/d | 200 | 100 | 100 |
| | TARGET RISK | LEVELS | | |
| Individual Excess Lifetime Cancer Risk | unitless | =adult residential | 1.0E-05 | 1.0E-05 |
| Hazard quotient | unitless | | 1.0 | 1.0 |